

# M13mp18 single-stranded DNA



1-800-632-7799  
info@neb.com  
www.neb.com



N4040S 015120914091

## N4040S

**10 µg** Lot: **0151209** Exp: **9/14**  
**250 µg/ml** Store at **-20°C**


**Description:** The single-stranded viral DNA is isolated from M13mp18. M13mp18 is a M13 *lac* phage vector which contains single HindIII, SphI, SbfI, PstI, SalI (AccI/ HincII), XbaI, BamHI, SmaI (XmaI), KpnI (Acc65I), SacI and EcoRI sites within the gene encoding β-Galactosidase. This DNA preparation is useful as a standard and has been tested as a template in the dideoxy-nucleotide termination method of sequencing DNA.

**Preparation:** M13mp18 phage is propagated in *E. coli* K12 JM101. The phage is purified by polyethylene glycol precipitation. The single-stranded DNA is then extracted with phenol.

Supplied in: 10 mM Tris-HCl (pH 8.0 @ 25°C), 1 mM EDTA.

### Reference:

1. Messing, J. (1983) *Methods Enzymol.* 101, 20.



CERTIFICATE OF ANALYSIS

# M13mp18 single-stranded DNA



1-800-632-7799  
info@neb.com  
www.neb.com



N4040S 015120914091

## N4040S

**10 µg** Lot: **0151209** Exp: **9/14**  
**250 µg/ml** Store at **-20°C**


**Description:** The single-stranded viral DNA is isolated from M13mp18. M13mp18 is a M13 *lac* phage vector which contains single HindIII, SphI, SbfI, PstI, SalI (AccI/ HincII), XbaI, BamHI, SmaI (XmaI), KpnI (Acc65I), SacI and EcoRI sites within the gene encoding β-Galactosidase. This DNA preparation is useful as a standard and has been tested as a template in the dideoxy-nucleotide termination method of sequencing DNA.

**Preparation:** M13mp18 phage is propagated in *E. coli* K12 JM101. The phage is purified by polyethylene glycol precipitation. The single-stranded DNA is then extracted with phenol.

Supplied in: 10 mM Tris-HCl (pH 8.0 @ 25°C), 1 mM EDTA.

### Reference:

1. Messing, J. (1983) *Methods Enzymol.* 101, 20.



CERTIFICATE OF ANALYSIS