

Endo H_f



1-800-632-7799
info@neb.com
www.neb.com



P0703S 018151017101

P0703S



100,000 units 1,000,000 U/ml Lot: 0181510

RECOMBINANT Store at -20°C Exp: 10/17

Description: Endo H_f is a recombinant protein fusion of Endoglycosidase H and maltose binding protein. Endo H_f cleaves the chitobiose core of high mannose and some hybrid oligosaccharides from N-linked glycoproteins (1) equally as well as Endo H.

New Reaction Buffer

Endo H_f



1-800-632-7799
info@neb.com
www.neb.com



P0703S 018151017101

P0703S




100,000 units 1,000,000 U/ml Lot: 0181510

RECOMBINANT Store at -20°C Exp: 10/17

Description: Endo H_f is a recombinant protein fusion of Endoglycosidase H and maltose binding protein. Endo H_f cleaves the chitobiose core of high mannose and some hybrid oligosaccharides from N-linked glycoproteins (1) equally as well as Endo H.

New Reaction Buffer

Specificity:



Source: Cloned from *Streptomyces plicatus* (2) and overexpressed in *E. coli* (3)

Applications:

- Removal of high mannose N-glycans from glycoproteins

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 5 mM Na₂EDTA.


Reagents Supplied with Enzyme:

10X Glycoprotein Denaturing Buffer:
5% SDS, **0.4 M DTT**

10X Glycobuffer 3:
[0.5 M Sodium Acetate (ph 6.0 @ 25°C)]

Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate.

Specificity:



Source: Cloned from *Streptomyces plicatus* (2) and overexpressed in *E. coli* (3)

Applications:

- Removal of high mannose N-glycans from glycoproteins

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 5 mM Na₂EDTA.

Reagents Supplied with Enzyme:

10X Glycoprotein Denaturing Buffer:
5% SDS, **0.4 M DTT**

10X Glycobuffer 3:
[0.5 M Sodium Acetate (ph 6.0 @ 25°C)]

Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate.

Reaction Conditions:

Typical reaction conditions are as follows:

- Combine 1–20 µg of glycoprotein, 1 µl of 10X Glycoprotein Denaturing Buffer and H₂O (if necessary) to make a 10 µl total reaction volume.
- Denature glycoprotein by heating reaction at 100°C for 10 minutes.
- Make a total reaction volume of 20 µl by adding 2 µl of 10X GlycoBuffer 3, H₂O and 1–5 µl Endo H.
- Incubate reaction at 37°C for 1 hour.

Note: Reactions may be scaled-up linearly to accommodate larger reaction volumes.

Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl (10 NEB units = 1 IUB milliunit).

Unit Definition Assay: 10 µg of RNase B are denatured with 1X Glycoprotein Denaturing Buffer at 100°C for 10 minutes. After the addition of 1X

Reaction Conditions:

Typical reaction conditions are as follows:

- Combine 1–20 µg of glycoprotein, 1 µl of 10X Glycoprotein Denaturing Buffer and H₂O (if necessary) to make a 10 µl total reaction volume.
- Denature glycoprotein by heating reaction at 100°C for 10 minutes.
- Make a total reaction volume of 20 µl by adding 2 µl of 10X GlycoBuffer 3, H₂O and 1–5 µl Endo H.
- Incubate reaction at 37°C for 1 hour.

Note: Reactions may be scaled-up linearly to accommodate larger reaction volumes.

Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl (10 NEB units = 1 IUB milliunit).

Unit Definition Assay: 10 µg of RNase B are denatured with 1X Glycoprotein Denaturing Buffer at 100°C for 10 minutes. After the addition of 1X

GlycoBuffer 3, two-fold dilutions of Endo H_f are added and the reaction mix is incubated for 1 hour at 37°C. Separation of reaction products are visualized by SDS-PAGE.

Specific Activity: ~232,000 units/mg.

Molecular Weight: 70,000 daltons.

Quality Assurance: No contaminating exoglycosidase or proteolytic activity could be detected.

Quality Controls

Glycosidase Assays: 5,000 units of Endo H were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10 µl reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

(See other side)

CERTIFICATE OF ANALYSIS

GlycoBuffer 3, two-fold dilutions of Endo H_f are added and the reaction mix is incubated for 1 hour at 37°C. Separation of reaction products are visualized by SDS-PAGE.

Specific Activity: ~232,000 units/mg.

Molecular Weight: 70,000 daltons.

Quality Assurance: No contaminating exoglycosidase or proteolytic activity could be detected.

Quality Controls

Glycosidase Assays: 5,000 units of Endo H were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10 µl reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

(See other side)

CERTIFICATE OF ANALYSIS

No other glycosidase activities were detected (ND) with the following substrates:

β-N-Acetyl-glucosaminidase:

GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC ND

α-Fucosidase:

Fucα1-2Galβ1-4Glc-AMC Galβ1-4
(Fucα1-3)GlcNAcβ1-3Galβ1-4Glc-AMC ND

β-Galactosidase:

Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC ND

α-Galactosidase:

Galα1-3Galβ1-4GlcNAc-AMC ND

α-Neuraminidase:

Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ
1-4Glc-AMC ND

α-Mannosidase:

Manα1-3Manβ1-4GlcNAc-AMC
Manα1-6Manα1-6(Manα1-3)Man-AMC ND

β-Glucosidase:

Glcβ1-4Glcβ1-4Glc-AMC ND

Page 2 (P0703)

No other glycosidase activities were detected (ND) with the following substrates:

β-N-Acetyl-glucosaminidase:

GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC ND

α-Fucosidase:

Fucα1-2Galβ1-4Glc-AMC Galβ1-4
(Fucα1-3)GlcNAcβ1-3Galβ1-4Glc-AMC ND

β-Galactosidase:

Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC ND

α-Galactosidase:

Galα1-3Galβ1-4GlcNAc-AMC ND

α-Neuraminidase:

Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ
1-4Glc-AMC ND

α-Mannosidase:

Manα1-3Manβ1-4GlcNAc-AMC
Manα1-6Manα1-6(Manα1-3)Man-AMC ND

β-Glucosidase:

Glcβ1-4Glcβ1-4Glc-AMC ND

Page 2 (P0703)

β-Xylosidase:

Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC ND

β-Mannosidase:

Manβ1-4Manβ1-4Man-AMC ND

Endo F₂, F₃:

Dansylated fibrinogen biantennary. ND

PNGase F:

Fluoresceinated fetuin triantennary. ND

Protease Assay: After incubation of 5,000 units of Endo H_i with 0.2 nmol of a standardized mixture of proteins, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

Notes On Use: Enzymatic activity is not affected by SDS.

To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

β-Xylosidase:

Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC ND

β-Mannosidase:

Manβ1-4Manβ1-4Man-AMC ND

Endo F₂, F₃:

Dansylated fibrinogen biantennary. ND

PNGase F:

Fluoresceinated fetuin triantennary. ND

Protease Assay: After incubation of 5,000 units of Endo H_i with 0.2 nmol of a standardized mixture of proteins, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

Notes On Use: Enzymatic activity is not affected by SDS.

To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

References:

1. Maley, F. et al. (1989) *Anal. Biochem.* 180, 195–204.
2. Robbins, P. et al. (1984) *J. Biol. Chem.* 259, 7577–7583.
3. Guan, C and Wong, S., New England Biolabs, Inc., unpublished results.

Companion Product Sold Separately:

RNase B
#P7817S 250 µg



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

References:

1. Maley, F. et al. (1989) *Anal. Biochem.* 180, 195–204.
2. Robbins, P. et al. (1984) *J. Biol. Chem.* 259, 7577–7583.
3. Guan, C and Wong, S., New England Biolabs, Inc., unpublished results.

Companion Product Sold Separately:

RNase B
#P7817S 250 µg



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.