



**β1-4 Galactosidase**



1-800-632-7799  
info@neb.com  
www.neb.com



P0730S 003150817081


**P0730S**



**400 units 8,000 U/ml Lot: 0031508**

**RECOMBINANT Store at -20°C Exp: 8/17**

**Description:** β1-4 Galactosidase is a highly specific exoglycosidase that catalyzes the hydrolysis of β1-4 linked D-galactopyranosyl residues from oligosaccharides.



**Detailed Specificity:** Specificity can vary depending on incubation time and branching structure.

**New Reaction Buffer**

**β1-4 Galactosidase**



1-800-632-7799  
info@neb.com  
www.neb.com



P0730S 003150817081


**P0730S**



**400 units 8,000 U/ml Lot: 0031508**


**RECOMBINANT Store at -20°C Exp: 8/17**

**Description:** β1-4 Galactosidase is a highly specific exoglycosidase that catalyzes the hydrolysis of β1-4 linked D-galactopyranosyl residues from oligosaccharides.




**Detailed Specificity:** Specificity can vary depending on incubation time and branching structure.

**New Reaction Buffer**



**Figure 1:** Detailed specificity of β1-4 Galactosidase. Reactions (A), (B) and (C) contained 2 units, 4 units and 8 units of β1-4 Galactosidase, respectively, and 1X GlycoBuffer 1 total reaction volume of 10 μl. Reactions were incubated at 37°C.



**Figure 1:** Detailed specificity of β1-4 Galactosidase. Reactions (A), (B) and (C) contained 2 units, 4 units and 8 units of β1-4 Galactosidase, respectively, and 1X GlycoBuffer 1 total reaction volume of 10 μl. Reactions were incubated at 37°C.

**Source:** Cloned from *Bacteroides fragilis* and expressed in *E. coli* (1).

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 1 mM Na<sub>2</sub>EDTA.

**Reagents Supplied with Enzyme:**  
10X GlycoBuffer 1

**Reaction Conditions:**  
1X GlycoBuffer 1:  
50 mM Sodium Acetate (pH 5.5 @ 25°C) and 5 mM CaCl<sub>2</sub>. Incubate at 37°C.

Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate.

**Unit Definition:** One unit is defined as the amount of enzyme required to cleave > 95% of the terminal, β-D-galactose from 1 nmol Galβ1-4GlcNAcβ1-3Galβ1-4Glc-7-amino-4-methylcoumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μl.

**Source:** Cloned from *Bacteroides fragilis* and expressed in *E. coli* (1).

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 1 mM Na<sub>2</sub>EDTA.

**Reagents Supplied with Enzyme:**  
10X GlycoBuffer 1

**Reaction Conditions:**  
1X GlycoBuffer 1:  
50 mM Sodium Acetate (pH 5.5 @ 25°C) and 5 mM CaCl<sub>2</sub>. Incubate at 37°C.

Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate.

**Unit Definition:** One unit is defined as the amount of enzyme required to cleave > 95% of the terminal, β-D-galactose from 1 nmol Galβ1-4GlcNAcβ1-3Galβ1-4Glc-7-amino-4-methylcoumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μl.

**Unit Definition Assay:** Two fold serial dilutions of β1-4 Galactosidase are incubated with 1 nmol AMC-labeled substrate in 1X GlycoBuffer 1, in a 10 μl reaction. The reaction mix is incubated for 1 hour at 37°C. Separation of reaction products are visualized via thin layer chromatography (2).

**Specific Activity:** 50,000 units/mg

**Molecular Weight:** 94,000 daltons.

**Quality Assurance:** No contaminating exoglycosidase or proteolytic activity could be detected.

**Quality Controls**  
**Glycosidase Assays:**  
32 units of β1-4 Galactosidase were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10 μl reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

(See other side)

CERTIFICATE OF ANALYSIS

**Unit Definition Assay:** Two fold serial dilutions of β1-4 Galactosidase are incubated with 1 nmol AMC-labeled substrate in 1X GlycoBuffer 1, in a 10 μl reaction. The reaction mix is incubated for 1 hour at 37°C. Separation of reaction products are visualized via thin layer chromatography (2).

**Specific Activity:** 50,000 units/mg

**Molecular Weight:** 94,000 daltons.

**Quality Assurance:** No contaminating exoglycosidase or proteolytic activity could be detected.

**Quality Controls**  
**Glycosidase Assays:**  
32 units of β1-4 Galactosidase were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10 μl reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

(See other side)

CERTIFICATE OF ANALYSIS

**Physical Purity:** Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

No other glycosidase activities were detected (ND) with the following substrates:

**$\beta$ -N-Acetylglucosaminidase:**

GlcNAc $\beta$ 1-4GlcNAc $\beta$ 1-4GlcNAc-AMC ND

**$\alpha$ -N-Acetylgalactosaminidase:**

GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Fucosidase:**

Fuc $\alpha$ 1-2Gal $\beta$ 1-4Glc-AMC ND

Gal $\beta$ 1-4 (Fuc $\alpha$ 1-3)GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Galactosidase:**

Gal $\alpha$ 1-3Gal $\beta$ 1-4Gal-AMC ND

Gal $\alpha$ 1-6Gal $\alpha$ 1-6Glc $\alpha$ 1-2Fru-AMC ND

**$\alpha$ -Neuraminidase:**

Neu5Ac $\alpha$ 2-3Gal $\beta$ 1-3GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Mannosidase:**

Man $\alpha$ 1-3Man $\beta$ 1-4GlcNAc-AMC ND

Man $\alpha$ 1-6Man $\alpha$ 1-6(Man $\alpha$ 1-3)Man-AMC ND

**$\beta$ -Glucosidase:**

Glc $\beta$ 1-4Glc $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Glucosidase:**

Glc $\alpha$ 1-6Glc $\alpha$ 1-4Glc-AMC ND

**$\beta$ -Xylosidase:**

Xyl $\beta$ 1-4Xyl $\beta$ 1-4Xyl $\beta$ 1-4Xyl-AMC ND

**$\beta$ -Mannosidase:**

Man $\beta$ 1-4Man $\beta$ 1-4Man-AMC ND

**Endo F<sub>1</sub>, F<sub>2</sub>, H:**

Dansylated invertase high mannose. ND

**Endo F<sub>2</sub>, F<sub>3</sub>:**

Dansylated fibrinogen biantennary. ND

**PNGase F:**

Fluoresceinated fetuin triantennary. ND

**Protease Assay:** After incubation of 112 units of  $\beta$ 1-4 Galactosidase with 0.2 nmol of a standard mixture of proteins in a 20  $\mu$ l reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

**Note:** Recommended storage temperature is -20°C.

**Heat Inactivation:** 65°C for 10 minutes.

**References:**

1. McLeod, E., New England Biolabs, Inc. unpublished results.
2. Wong-Madden, S.T. and Landry, D. (1995) *Glycobiology* 5, 19–28.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

**Physical Purity:** Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

No other glycosidase activities were detected (ND) with the following substrates:

**$\beta$ -N-Acetylglucosaminidase:**

GlcNAc $\beta$ 1-4GlcNAc $\beta$ 1-4GlcNAc-AMC ND

**$\alpha$ -N-Acetylgalactosaminidase:**

GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Fucosidase:**

Fuc $\alpha$ 1-2Gal $\beta$ 1-4Glc-AMC ND

Gal $\beta$ 1-4 (Fuc $\alpha$ 1-3)GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Galactosidase:**

Gal $\alpha$ 1-3Gal $\beta$ 1-4Gal-AMC ND

Gal $\alpha$ 1-6Gal $\alpha$ 1-6Glc $\alpha$ 1-2Fru-AMC ND

**$\alpha$ -Neuraminidase:**

Neu5Ac $\alpha$ 2-3Gal $\beta$ 1-3GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Mannosidase:**

Man $\alpha$ 1-3Man $\beta$ 1-4GlcNAc-AMC ND

Man $\alpha$ 1-6Man $\alpha$ 1-6(Man $\alpha$ 1-3)Man-AMC ND

**$\beta$ -Glucosidase:**

Glc $\beta$ 1-4Glc $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Glucosidase:**

Glc $\alpha$ 1-6Glc $\alpha$ 1-4Glc-AMC ND

**$\beta$ -Xylosidase:**

Xyl $\beta$ 1-4Xyl $\beta$ 1-4Xyl $\beta$ 1-4Xyl-AMC ND

**$\beta$ -Mannosidase:**

Man $\beta$ 1-4Man $\beta$ 1-4Man-AMC ND

**Endo F<sub>1</sub>, F<sub>2</sub>, H:**

Dansylated invertase high mannose. ND

**Endo F<sub>2</sub>, F<sub>3</sub>:**

Dansylated fibrinogen biantennary. ND

**PNGase F:**

Fluoresceinated fetuin triantennary. ND

**Protease Assay:** After incubation of 112 units of  $\beta$ 1-4 Galactosidase with 0.2 nmol of a standard mixture of proteins in a 20  $\mu$ l reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

**Note:** Recommended storage temperature is -20°C.

**Heat Inactivation:** 65°C for 10 minutes.

**References:**

1. McLeod, E., New England Biolabs, Inc. unpublished results.
2. Wong-Madden, S.T. and Landry, D. (1995) *Glycobiology* 5, 19–28.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.