

## O-Glycosidase



1-800-632-7799  
info@neb.com  
www.neb.com



P0733S 001150417041

# P0733S



2,000,000 units 40,000,000 U/ml Lot: 0011504

RECOMBINANT Store at -20°C Exp: 4/17

**Description:** *O*-Glycosidase, also known as Endo- $\alpha$ -*N*-Acetylgalactosaminidase, catalyzes the removal of Core 1 and Core 3 *O*-linked disaccharides from glycoproteins.

### Specificity:

A. Core 1



B. Core 3



## O-Glycosidase



1-800-632-7799  
info@neb.com  
www.neb.com



P0733S 001150417041

# P0733S



2,000,000 units 40,000,000 U/ml Lot: 0011504

RECOMBINANT Store at -20°C Exp: 4/17

**Description:** *O*-Glycosidase, also known as Endo- $\alpha$ -*N*-Acetylgalactosaminidase, catalyzes the removal of Core 1 and Core 3 *O*-linked disaccharides from glycoproteins.

### Specificity:

A. Core 1



B. Core 3



**Source:** Cloned from *Enterococcus faecalis* and expressed in *E. coli* (1).

### Applications:

- Removal of Core 1 & Core 3 *O*-linked disaccharide glycans from glycoproteins

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 1 mM Na<sub>2</sub>EDTA.

### Reagents Supplied with Enzyme:

10X Glycoprotein Denaturing Buffer

10X GlycoBuffer 2

10% NP-40

**Unit Definition:** One unit is defined as the amount of enzyme required to remove 0.68 nmol of *O*-linked disaccharide from 5 mg of neuraminidase digested, non-denatured fetuin (2) in 1 hour at 37°C in a total reaction volume of 100  $\mu$ l (1 unit of both *O*-Glycosidase and PNGase F will remove equivalent molar amounts of *O*-linked disaccharides and *N*-linked oligosaccharides, respectively).

**Source:** Cloned from *Enterococcus faecalis* and expressed in *E. coli* (1).

### Applications:

- Removal of Core 1 & Core 3 *O*-linked disaccharide glycans from glycoproteins

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 1 mM Na<sub>2</sub>EDTA.

### Reagents Supplied with Enzyme:

10X Glycoprotein Denaturing Buffer

10X GlycoBuffer 2

10% NP-40

**Unit Definition:** One unit is defined as the amount of enzyme required to remove 0.68 nmol of *O*-linked disaccharide from 5 mg of neuraminidase digested, non-denatured fetuin (2) in 1 hour at 37°C in a total reaction volume of 100  $\mu$ l (1 unit of both *O*-Glycosidase and PNGase F will remove equivalent molar amounts of *O*-linked disaccharides and *N*-linked oligosaccharides, respectively).

**Non-Denaturing Unit Definition Assay:** Two fold serial dilutions of *O*-Glycosidase are added to a reaction mixture of 5 mg of neuraminidase digested fetuin with 1X GlycoBuffer 2. The reaction mix is then incubated at 37°C for 1 hour. *O*-linked disaccharide carbohydrates are determined by the Morgan and Elson Assay (2).

**Note:** Under denaturing conditions the enzyme activity is increased two-fold. This observation is substrate dependent.

**Specific Activity:** 53,000,000 units/mg.

**Molecular Weight:** 147,000 daltons.

**Quality Assurance:** No contaminating exoglycosidase or proteolytic activity could be detected.

### Quality Controls

#### Glycosidase Assays:

200,000 units of *O*-Glycosidase were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10  $\mu$ l reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

**Non-Denaturing Unit Definition Assay:** Two fold serial dilutions of *O*-Glycosidase are added to a reaction mixture of 5 mg of neuraminidase digested fetuin with 1X GlycoBuffer 2. The reaction mix is then incubated at 37°C for 1 hour. *O*-linked disaccharide carbohydrates are determined by the Morgan and Elson Assay (2).

**Note:** Under denaturing conditions the enzyme activity is increased two-fold. This observation is substrate dependent.

**Specific Activity:** 53,000,000 units/mg.

**Molecular Weight:** 147,000 daltons.

**Quality Assurance:** No contaminating exoglycosidase or proteolytic activity could be detected.

### Quality Controls

#### Glycosidase Assays:

200,000 units of *O*-Glycosidase were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10  $\mu$ l reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

No other glycosidase activities were detected (ND) with the following substrates:

**$\beta$ -*N*-Acetylgalactosaminidase:**  
GalNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -*N*-Acetylgalactosaminidase:**  
GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Fucosidase:**  
Fuc $\alpha$ 1-2Gal $\beta$ 1-4Glc-AMC ND  
Gal $\beta$ 1-4 (Fuc $\alpha$ 1-3)GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\beta$ -Galactosidase:**  
Gal $\beta$ 1-3GlcNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC ND  
Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Galactosidase:**  
Gal $\alpha$ 1-3Gal $\beta$ 1-4Gal-AMC ND  
Gal $\alpha$ 1-6Gal $\alpha$ 1-6Glc $\alpha$ 1-2Fru-AMC ND

(see other side)

CERTIFICATE OF ANALYSIS

No other glycosidase activities were detected (ND) with the following substrates:

**$\beta$ -*N*-Acetylgalactosaminidase:**  
GalNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -*N*-Acetylgalactosaminidase:**  
GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Fucosidase:**  
Fuc $\alpha$ 1-2Gal $\beta$ 1-4Glc-AMC ND  
Gal $\beta$ 1-4 (Fuc $\alpha$ 1-3)GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\beta$ -Galactosidase:**  
Gal $\beta$ 1-3GlcNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC ND  
Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Galactosidase:**  
Gal $\alpha$ 1-3Gal $\beta$ 1-4Gal-AMC ND  
Gal $\alpha$ 1-6Gal $\alpha$ 1-6Glc $\alpha$ 1-2Fru-AMC ND

(see other side)

CERTIFICATE OF ANALYSIS

**$\alpha$ -Neuraminidase:**  
Neu5Ac $\alpha$ 2-3Gal $\beta$ 1-3GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Mannosidase:**  
Man $\alpha$ 1-3Man $\beta$ 1-4GlcNAc-AMC ND  
Man $\alpha$ 1-6Man $\alpha$ 1-6(Man $\alpha$ 1-3)Man-AMC ND

**$\beta$ -Glucosidase:**  
Glc $\beta$ 1-4Glc $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Glucosidase:**  
Glc $\alpha$ 1-6Glc $\alpha$ 1-4Glc-AMC ND

**$\beta$ -Xylosidase:**  
Xyl $\beta$ 1-4Xyl $\beta$ 1-4Xyl $\beta$ 1-4Xyl-AMC ND

**$\beta$ -Mannosidase:**  
Man $\beta$ 1-4Man $\beta$ 1-4Man-AMC ND

**Endo F<sub>1</sub>, F<sub>2</sub>, H:**  
Dansylated invertase high mannose. ND

**Endo F<sub>2</sub>, F<sub>3</sub>:**  
Dansylated fibrinogen biantennary. ND

Page 2 (P0733)

**$\alpha$ -Neuraminidase:**  
Neu5Ac $\alpha$ 2-3Gal $\beta$ 1-3GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Mannosidase:**  
Man $\alpha$ 1-3Man $\beta$ 1-4GlcNAc-AMC ND  
Man $\alpha$ 1-6Man $\alpha$ 1-6(Man $\alpha$ 1-3)Man-AMC ND

**$\beta$ -Glucosidase:**  
Glc $\beta$ 1-4Glc $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Glucosidase:**  
Glc $\alpha$ 1-6Glc $\alpha$ 1-4Glc-AMC ND

**$\beta$ -Xylosidase:**  
Xyl $\beta$ 1-4Xyl $\beta$ 1-4Xyl $\beta$ 1-4Xyl-AMC ND

**$\beta$ -Mannosidase:**  
Man $\beta$ 1-4Man $\beta$ 1-4Man-AMC ND

**Endo F<sub>1</sub>, F<sub>2</sub>, H:**  
Dansylated invertase high mannose. ND

**Endo F<sub>2</sub>, F<sub>3</sub>:**  
Dansylated fibrinogen biantennary. ND

Page 2 (P0733)

**PNGase F:**  
Fluoresceinated fetuin triantennary. ND

**Protease Assay:** After incubation of 1,400,000 units of *O*-Glycosidase with 0.2 nmol of a standard mixture of proteins in a 20  $\mu$ l reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

**Protocol:** Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate. Typical reaction conditions are as follows:

1. Combine 10–20  $\mu$ g of glycoprotein, 1  $\mu$ l of 10X Glycoprotein Denaturing Buffer and H<sub>2</sub>O (if necessary) to make a 10  $\mu$ l total reaction volume.
2. Denature glycoprotein by heating reaction at 100°C for 10 minutes.
3. Make a total reaction volume of 20  $\mu$ l by adding 2  $\mu$ l 10X GlycoBuffer 2, 2  $\mu$ l 10% NP40, 2  $\mu$ l Neuraminidase, H<sub>2</sub>O and 1–5  $\mu$ l *O*-Glycosidase.
4. Incubate reaction at 37°C for 1–4 hours.

**PNGase F:**  
Fluoresceinated fetuin triantennary. ND

**Protease Assay:** After incubation of 1,400,000 units of *O*-Glycosidase with 0.2 nmol of a standard mixture of proteins in a 20  $\mu$ l reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

**Protocol:** Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate. Typical reaction conditions are as follows:

1. Combine 10–20  $\mu$ g of glycoprotein, 1  $\mu$ l of 10X Glycoprotein Denaturing Buffer and H<sub>2</sub>O (if necessary) to make a 10  $\mu$ l total reaction volume.
2. Denature glycoprotein by heating reaction at 100°C for 10 minutes.
3. Make a total reaction volume of 20  $\mu$ l by adding 2  $\mu$ l 10X GlycoBuffer 2, 2  $\mu$ l 10% NP40, 2  $\mu$ l Neuraminidase, H<sub>2</sub>O and 1–5  $\mu$ l *O*-Glycosidase.
4. Incubate reaction at 37°C for 1–4 hours.

**Notes on Use:** Since *O*-Glycosidase is inhibited by SDS, it is essential to have NP-40 in the reaction mixture. It is not known why this non-ionic detergent counteracts the SDS inhibition at the present time. Double digest with Endo H must have NP-40 present (NP-40 does not inhibit Endo H).

To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

Recommended storage temperature is –20°C.

**Heat Inactivation:** 65°C for 10 minutes.

**Notes on Use:** Since *O*-Glycosidase is inhibited by SDS, it is essential to have NP-40 in the reaction mixture. It is not known why this non-ionic detergent counteracts the SDS inhibition at the present time. Double digest with Endo H must have NP-40 present (NP-40 does not inhibit Endo H).

To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

Recommended storage temperature is –20°C.

**Heat Inactivation:** 65°C for 10 minutes.

## References:

1. Koutsoulis, D., Landry, D. and Guthrie, E.P. (2008) *Glycobiology* 18, 799–805.
2. Morgan, W.T.J. and Elson, L.A. (1934) *Biochem. J.* 28, 988–995.



NEW ENGLAND BIOLABS® and REMOVE-IT® are registered trademarks of New England Biolabs, Inc.

## References:

1. Koutsoulis, D., Landry, D. and Guthrie, E.P. (2008) *Glycobiology* 18, 799–805.
2. Morgan, W.T.J. and Elson, L.A. (1934) *Biochem. J.* 28, 988–995.



NEW ENGLAND BIOLABS® and REMOVE-IT® are registered trademarks of New England Biolabs, Inc.