

## O-Glycosidase



1-800-632-7799  
info@neb.com  
www.neb.com



P0733S 003140816081

# P0733S



**2,000,000 units 40,000,000 U/ml Lot: 0031408**  
**RECOMBINANT Store at -20°C Exp: 8/16**

**Description:** *O*-Glycosidase, also known as Endo- $\alpha$ -*N*-Acetylgalactosaminidase, catalyzes the removal of Core 1 and Core 3 *O*-linked disaccharides from glycoproteins.

### Specificity:

A. Core 1



B. Core 3



## O-Glycosidase



1-800-632-7799  
info@neb.com  
www.neb.com



P0733S 003140816081

# P0733S



**2,000,000 units 40,000,000 U/ml Lot: 0031408**  
**RECOMBINANT Store at -20°C Exp: 8/16**

**Description:** *O*-Glycosidase, also known as Endo- $\alpha$ -*N*-Acetylgalactosaminidase, catalyzes the removal of Core 1 and Core 3 *O*-linked disaccharides from glycoproteins.

### Specificity:

A. Core 1



B. Core 3



**Source:** Cloned from *Enterococcus faecalis* and expressed in *E. coli* (1).

### Applications:

- Removal of Core 1 & Core 3 *O*-linked disaccharide glycans from glycoproteins

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 1 mM Na<sub>2</sub>EDTA.

### Reagents Supplied with Enzyme:

10X Glycoprotein Denaturing Buffer  
10X G7 Reaction Buffer  
10% NP-40

**Unit Definition:** One unit is defined as the amount of enzyme required to remove 0.68 nmol of *O*-linked disaccharide from 5 mg of neuraminidase digested, non-denatured fetuin (2) in 1 hour at 37°C in a total reaction volume of 100  $\mu$ l (1 unit of both *O*-Glycosidase and PNGase F will remove equivalent molar amounts of *O*-linked disaccharides and *N*-linked oligosaccharides, respectively).

**Source:** Cloned from *Enterococcus faecalis* and expressed in *E. coli* (1).

### Applications:

- Removal of Core 1 & Core 3 *O*-linked disaccharide glycans from glycoproteins

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 1 mM Na<sub>2</sub>EDTA.

### Reagents Supplied with Enzyme:

10X Glycoprotein Denaturing Buffer  
10X G7 Reaction Buffer  
10% NP-40

**Unit Definition:** One unit is defined as the amount of enzyme required to remove 0.68 nmol of *O*-linked disaccharide from 5 mg of neuraminidase digested, non-denatured fetuin (2) in 1 hour at 37°C in a total reaction volume of 100  $\mu$ l (1 unit of both *O*-Glycosidase and PNGase F will remove equivalent molar amounts of *O*-linked disaccharides and *N*-linked oligosaccharides, respectively).

**Non-Denaturing Unit Definition Assay:** Two fold serial dilutions of *O*-Glycosidase are added to a reaction mixture of 5 mg of neuraminidase digested fetuin with 1X G7 Reaction Buffer. The reaction mix is then incubated at 37°C for 1 hour. *O*-linked disaccharide carbohydrates are determined by the Morgan and Elson Assay (2).

**Note:** Under denaturing conditions the enzyme activity is increased two-fold. This observation is substrate dependent.

**Specific Activity:** 53,000,000 units/mg.

**Molecular Weight:** 147,000 daltons.

**Quality Assurance:** No contaminating exoglycosidase or proteolytic activity could be detected.

**Non-Denaturing Unit Definition Assay:** Two fold serial dilutions of *O*-Glycosidase are added to a reaction mixture of 5 mg of neuraminidase digested fetuin with 1X G7 Reaction Buffer. The reaction mix is then incubated at 37°C for 1 hour. *O*-linked disaccharide carbohydrates are determined by the Morgan and Elson Assay (2).

**Note:** Under denaturing conditions the enzyme activity is increased two-fold. This observation is substrate dependent.

**Specific Activity:** 53,000,000 units/mg.

**Molecular Weight:** 147,000 daltons.

**Quality Assurance:** No contaminating exoglycosidase or proteolytic activity could be detected.

### Quality Controls

#### Glycosidase Assays:

200,000 units of *O*-Glycosidase were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10  $\mu$ l reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

No other glycosidase activities were detected (ND) with the following substrates:

**$\beta$ -*N*-Acetylgalactosaminidase:**  
GalNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -*N*-Acetylgalactosaminidase:**  
GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Fucosidase:**  
Fuc $\alpha$ 1-2Gal $\beta$ 1-4Glc-AMC ND  
Gal $\beta$ 1-4 (Fuc $\alpha$ 1-3)GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

(see other side)

CERTIFICATE OF ANALYSIS

### Quality Controls

#### Glycosidase Assays:

200,000 units of *O*-Glycosidase were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10  $\mu$ l reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

No other glycosidase activities were detected (ND) with the following substrates:

**$\beta$ -*N*-Acetylgalactosaminidase:**  
GalNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -*N*-Acetylgalactosaminidase:**  
GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4Glc-AMC ND

**$\alpha$ -Fucosidase:**  
Fuc $\alpha$ 1-2Gal $\beta$ 1-4Glc-AMC ND  
Gal $\beta$ 1-4 (Fuc $\alpha$ 1-3)GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC ND

(see other side)

CERTIFICATE OF ANALYSIS

<b>β-Galactosidase:</b> Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC	ND
Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC	ND
<b>α-Galactosidase:</b> Galα1-3Galβ1-4Gal-AMC	ND
Galα1-6Galα1-6Glcα1-2Fru-AMC	ND
<b>α-Neuraminidase:</b> Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC	ND
<b>α-Mannosidase:</b> Manα1-3Manβ1-4GlcNAc-AMC	ND
Manα1-6Manα1-6(Manα1-3)Man-AMC	ND
<b>β-Glucosidase:</b> Glcβ1-4Glcβ1-4Glc-AMC	ND
<b>α-Glucosidase:</b> Glcα1-6Glcα1-4Glc-AMC	ND
<b>β-Xylosidase:</b> Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC	ND

Page 2 (P0733)

<b>β-Galactosidase:</b> Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC	ND
Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC	ND
<b>α-Galactosidase:</b> Galα1-3Galβ1-4Gal-AMC	ND
Galα1-6Galα1-6Glcα1-2Fru-AMC	ND
<b>α-Neuraminidase:</b> Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC	ND
<b>α-Mannosidase:</b> Manα1-3Manβ1-4GlcNAc-AMC	ND
Manα1-6Manα1-6(Manα1-3)Man-AMC	ND
<b>β-Glucosidase:</b> Glcβ1-4Glcβ1-4Glc-AMC	ND
<b>α-Glucosidase:</b> Glcα1-6Glcα1-4Glc-AMC	ND
<b>β-Xylosidase:</b> Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC	ND

Page 2 (P0733)

<b>β-Mannosidase:</b> Manβ1-4Manβ1-4Man-AMC	ND
<b>Endo F<sub>1</sub>, F<sub>2</sub>, H:</b> Dansylated invertase high mannose.	ND
<b>Endo F<sub>2</sub>, F<sub>3</sub>:</b> Dansylated fibrinogen biantennary.	ND
<b>PNGase F:</b> Fluoresceinated fetuin triantennary.	ND

**Protease Assay:** After incubation of 1,400,000 units of *O*-Glycosidase with 0.2 nmol of a standard mixture of proteins in a 20 μl reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

<b>β-Mannosidase:</b> Manβ1-4Manβ1-4Man-AMC	ND
<b>Endo F<sub>1</sub>, F<sub>2</sub>, H:</b> Dansylated invertase high mannose.	ND
<b>Endo F<sub>2</sub>, F<sub>3</sub>:</b> Dansylated fibrinogen biantennary.	ND
<b>PNGase F:</b> Fluoresceinated fetuin triantennary.	ND

**Protease Assay:** After incubation of 1,400,000 units of *O*-Glycosidase with 0.2 nmol of a standard mixture of proteins in a 20 μl reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

**Protocol:** Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate. Typical reaction conditions are as follows:

1. Combine 10–20 μg of glycoprotein, 1 μl of 10X Glycoprotein Denaturing Buffer and H<sub>2</sub>O (if necessary) to make a 10 μl total reaction volume.
2. Denature glycoprotein by heating reaction at 100°C for 10 minutes.
3. Make a total reaction volume of 20 μl by adding 2 μl 10X G7 Reaction Buffer, 2 μl 10% NP40, 2 μl Neuraminidase, H<sub>2</sub>O and 1–5 μl *O*-Glycosidase.
4. Incubate reaction at 37°C for 1–4 hours.

**Protocol:** Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate. Typical reaction conditions are as follows:

1. Combine 10–20 μg of glycoprotein, 1 μl of 10X Glycoprotein Denaturing Buffer and H<sub>2</sub>O (if necessary) to make a 10 μl total reaction volume.
2. Denature glycoprotein by heating reaction at 100°C for 10 minutes.
3. Make a total reaction volume of 20 μl by adding 2 μl 10X G7 Reaction Buffer, 2 μl 10% NP40, 2 μl Neuraminidase, H<sub>2</sub>O and 1–5 μl *O*-Glycosidase.
4. Incubate reaction at 37°C for 1–4 hours.

**Notes on Use:** Since *O*-Glycosidase is inhibited by SDS, it is essential to have NP-40 in the reaction mixture. It is not known why this non-ionic detergent counteracts the SDS inhibition at the present time. Double digest with Endo H must have NP-40 present (NP-40 does not inhibit Endo H).

To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

Recommended storage temperature is –20°C.

**Heat Inactivation:** 65°C for 10 minutes.

#### References:

1. Koutsioulis, D., Landry, D. and Guthrie, E.P. (2008) *Glycobiology* 18, 799–805.
2. Morgan, W.T.J. and Elson, L.A. (1934) *Biochem. J.* 28, 988–995.

**Notes on Use:** Since *O*-Glycosidase is inhibited by SDS, it is essential to have NP-40 in the reaction mixture. It is not known why this non-ionic detergent counteracts the SDS inhibition at the present time. Double digest with Endo H must have NP-40 present (NP-40 does not inhibit Endo H).

To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

Recommended storage temperature is –20°C.

**Heat Inactivation:** 65°C for 10 minutes.

#### References:

1. Koutsioulis, D., Landry, D. and Guthrie, E.P. (2008) *Glycobiology* 18, 799–805.
2. Morgan, W.T.J. and Elson, L.A. (1934) *Biochem. J.* 28, 988–995.