

Casein Kinase II (CK2)



1-800-632-7799
info@neb.com
www.neb.com



P6010S 016150416041

P6010S



10,000 units 500,000 U/ml Lot: 0161504

RECOMBINANT Store at -80°C Exp: 4/16

Description: Casein Kinase II (CK2) is a constitutively active serine/threonine protein kinase composed of two 44 kDa catalytic α -subunits and two 26 kDa regulatory β -subunits in an $\alpha_2\beta_2$ configuration to form stable heterotetramers. CK2 holoenzyme undergoes autophosphorylation at two serine residues (S2/S3) of its β -subunit. Recently it has been shown that CK2 α -subunits undergo intermolecular tyrosine-autophosphorylation at Y182, which may represent a specific regulatory

New Reaction Buffer

Casein Kinase II (CK2)



1-800-632-7799
info@neb.com
www.neb.com



P6010S 016150416041

P6010S



10,000 units 500,000 U/ml Lot: 0161504

RECOMBINANT Store at -80°C Exp: 4/16

Description: Casein Kinase II (CK2) is a constitutively active serine/threonine protein kinase composed of two 44 kDa catalytic α -subunits and two 26 kDa regulatory β -subunits in an $\alpha_2\beta_2$ configuration to form stable heterotetramers. CK2 holoenzyme undergoes autophosphorylation at two serine residues (S2/S3) of its β -subunit. Recently it has been shown that CK2 α -subunits undergo intermolecular tyrosine-autophosphorylation at Y182, which may represent a specific regulatory

New Reaction Buffer

mechanism. Also, CK2 is able to phosphorylate, under special circumstances, tyrosyl residues in proteins. CK2 is implicated in a variety of cellular functions (1,2).

Recognition Determinants: The CK2 substrate specificity is invariably determined by multiple acidic residues located at positions between -2 and +5 relative to the target amino acid (mostly Ser and rarely Thr). The general recognition motif for phosphorylation by CK2 is **SXXE/D**, although **SXE/D** and **S/D**, and variations of these sequences are also phosphorylated. Polyanionic compounds, like heparin, inhibit CK2 activity with a K_i of 1.4 nM (4,5).

Source: Isolated from a strain of *E. coli* expressing both α and β CK2 subunits derived from a human glioblastoma cDNA library (kindly provided by Dr. D. Marshak) (3).

Supplied in: 350 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C), 1 mM Na₂EDTA, 2 mM DTT and 0.1% Triton X-100.

mechanism. Also, CK2 is able to phosphorylate, under special circumstances, tyrosyl residues in proteins. CK2 is implicated in a variety of cellular functions (1,2).

Recognition Determinants: The CK2 substrate specificity is invariably determined by multiple acidic residues located at positions between -2 and +5 relative to the target amino acid (mostly Ser and rarely Thr). The general recognition motif for phosphorylation by CK2 is **SXXE/D**, although **SXE/D** and **S/D**, and variations of these sequences are also phosphorylated. Polyanionic compounds, like heparin, inhibit CK2 activity with a K_i of 1.4 nM (4,5).

Source: Isolated from a strain of *E. coli* expressing both α and β CK2 subunits derived from a human glioblastoma cDNA library (kindly provided by Dr. D. Marshak) (3).

Supplied in: 350 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C), 1 mM Na₂EDTA, 2 mM DTT and 0.1% Triton X-100.

Reagents Supplied with Enzyme:
10X NEBuffer for Protein Kinases

Reaction Conditions: 1X NEBuffer for Protein Kinases, supplement with 200 μ M ATP and gamma-labeled ATP to a final specific activity of 100–500 μ Ci/ μ mol. (CK2 will also accept GTP as a phosphoryl donor in place of ATP). **Incubate at 30°C.**

1X NEBuffer for Protein Kinases:

50 mM Tris-HCl
10 mM MgCl₂
0.1 mM EDTA
2 mM DTT
0.01% Brij 35
pH 7.5 @ 25°C

Note that optimal incubation times and enzyme concentrations must be determined empirically for each particular substrate.

Unit Definition: One unit is defined as the amount of CK2 required to catalyze the transfer of 1 pmol of phosphate to CK2 Peptide Substrate, RRRADSDDDDD (100 μ M, NEB #P6012), in 1 minute at 30°C in a total reaction volume of 25 μ l (4,5).

Reagents Supplied with Enzyme:
10X NEBuffer for Protein Kinases

Reaction Conditions: 1X NEBuffer for Protein Kinases, supplement with 200 μ M ATP and gamma-labeled ATP to a final specific activity of 100–500 μ Ci/ μ mol. (CK2 will also accept GTP as a phosphoryl donor in place of ATP). **Incubate at 30°C.**

1X NEBuffer for Protein Kinases:

50 mM Tris-HCl
10 mM MgCl₂
0.1 mM EDTA
2 mM DTT
0.01% Brij 35
pH 7.5 @ 25°C

Note that optimal incubation times and enzyme concentrations must be determined empirically for each particular substrate.

Unit Definition: One unit is defined as the amount of CK2 required to catalyze the transfer of 1 pmol of phosphate to CK2 Peptide Substrate, RRRADSDDDDD (100 μ M, NEB #P6012), in 1 minute at 30°C in a total reaction volume of 25 μ l (4,5).

Specific Activity: ~ 859,000 units/mg.

Molecular Weight: α -subunit (45 kDa), β -subunit (25 kDa). The apparent molecular weight of the α -subunit estimated by SDS-PAGE is about 42 kDa.

Quality Assurance: CK2 contains no detectable protease or phosphatase activities.

Quality Control Assays

Protease Activity: After incubation of 5,000 units of Casein Kinase II (CK2) with a standard mixture of proteins for 2 hours at 30°C, no proteolytic activity could be detected by SDS-PAGE analysis.

Phosphatase Activity: After incubation of 5,000 units of Casein Kinase II (CK2) with 50 mM *p*-nitrophenyl phosphate for 2 hours at 30°C, no phosphatase activity could be detected by spectrophotometric analysis.

(See other side)

CERTIFICATE OF ANALYSIS

Specific Activity: ~ 859,000 units/mg.

Molecular Weight: α -subunit (45 kDa), β -subunit (25 kDa). The apparent molecular weight of the α -subunit estimated by SDS-PAGE is about 42 kDa.

Quality Assurance: CK2 contains no detectable protease or phosphatase activities.

Quality Control Assays

Protease Activity: After incubation of 5,000 units of Casein Kinase II (CK2) with a standard mixture of proteins for 2 hours at 30°C, no proteolytic activity could be detected by SDS-PAGE analysis.

Phosphatase Activity: After incubation of 5,000 units of Casein Kinase II (CK2) with 50 mM *p*-nitrophenyl phosphate for 2 hours at 30°C, no phosphatase activity could be detected by spectrophotometric analysis.

(See other side)

CERTIFICATE OF ANALYSIS

References:

1. Donella-Deana, A. et al. (2001) *Biochem J.* 357, 563–567.
2. Marin, O. et al. (1999) *J. Biol. Chem.* 274, 29260–29265.
3. Chester, N. and Marshak, D.R. (1993) *Anal. Biochem.* 209, 284–290.
4. Marin, O. et al. (1994) *BBRC*, 198, 898–905.
5. Sarno, S. et al. (1996) *J. Biol. Chem.* 271, 10595–10601.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

References:

1. Donella-Deana, A. et al. (2001) *Biochem J.* 357, 563–567.
2. Marin, O. et al. (1999) *J. Biol. Chem.* 274, 29260–29265.
3. Chester, N. and Marshak, D.R. (1993) *Anal. Biochem.* 209, 284–290.
4. Marin, O. et al. (1994) *BBRC*, 198, 898–905.
5. Sarno, S. et al. (1996) *J. Biol. Chem.* 271, 10595–10601.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.