

Glycogen Synthase Kinase 3 (GSK-3)



1-800-632-7799
info@neb.com
www.neb.com



P6040S 029160217021

P6040S



10,000 units 500,000 U/ml Lot: 0291602
RECOMBINANT Store at -20°C Exp: 2/17

Description: Glycogen Synthase Kinase 3 (GSK-3) is a serine/threonine protein kinase and one of several protein kinases, which phosphorylate glycogen synthase. It is also called Factor A (F_a) for its ability to activate the MgATP-dependent form of the protein phosphatase PP1 called F_c (1–4). Recent studies demonstrate that GSK-3 can autophosphorylate Ser, Thr and Tyr. Ser/Thr phosphorylation causes inactivation, and

New Reaction Buffer

Tyr phosphorylation results in increased activity (Y216 for GSK-3 β). GSK-3 expressed in *E. coli* or insect cells is extensively phosphorylated on Tyr. Molecules lacking phosphate at this position can autophosphorylate after incubation with Mg²⁺ and ATP. GSK-3 phosphorylates several exogenous substrates, but not on Tyr residues (5,6).

Recognition Determinants: The substrate specificity of GSK-3 is unique and substrate dependent. For some substrates, prior phosphorylation of the substrate to form the motif S/TXXXpS/pT is a strict requirement whereas in other substrates, no previous phosphorylation is needed. In either case, many of the GSK-3 sites have Pro residues close to the modified Ser or Thr (5,7).

Source: Isolated from a strain of *E. coli* that carries a clone expressing GSK-3 β derived from a rabbit skeletal muscle cDNA library (kindly provided by Dr. P.J. Roach) (5).

Supplied in: 50 mM NaCl, 30 mM Tris-HCl (pH 7.5 @ 25°C), 1.0 mM EDTA, 5 mM DTT, 0.03% Brij and 50% glycerol.

Tyr phosphorylation results in increased activity (Y216 for GSK-3 β). GSK-3 expressed in *E. coli* or insect cells is extensively phosphorylated on Tyr. Molecules lacking phosphate at this position can autophosphorylate after incubation with Mg²⁺ and ATP. GSK-3 phosphorylates several exogenous substrates, but not on Tyr residues (5,6).

Recognition Determinants: The substrate specificity of GSK-3 is unique and substrate dependent. For some substrates, prior phosphorylation of the substrate to form the motif S/TXXXpS/pT is a strict requirement whereas in other substrates, no previous phosphorylation is needed. In either case, many of the GSK-3 sites have Pro residues close to the modified Ser or Thr (5,7).

Source: Isolated from a strain of *E. coli* that carries a clone expressing GSK-3 β derived from a rabbit skeletal muscle cDNA library (kindly provided by Dr. P.J. Roach) (5).

Supplied in: 50 mM NaCl, 30 mM Tris-HCl (pH 7.5 @ 25°C), 1.0 mM EDTA, 5 mM DTT, 0.03% Brij and 50% glycerol.

Reagents Supplied with Enzyme:
10X NEBuffer for Protein Kinases

Reaction Conditions: 1X NEBuffer for Protein Kinases, supplemented with 200 μ M ATP and gamma-labeled ATP to a final specific activity of 100–500 μ Ci/ μ mol. **Incubate at 30°C.**

1X NEBuffer for Protein Kinases:

50 mM Tris-HCl
10 mM MgCl₂
0.1 mM EDTA
2 mM DTT
0.01% Brij 35
pH 7.5 @ 25°C

Note that optimal incubation times and enzyme concentrations must be determined empirically for each particular substrate.

Unit Definition: One unit is defined as the amount of GSK-3 required to catalyze the transfer of 1 pmol of phosphate to CREB Phosphopeptide, KRREILSRPPpSYR (400 μ M, NEB #P6041), in 1 minute at 30°C in a total reaction volume of 25 μ l.

Specific Activity: ~ 5,000,000 units/mg.

Reagents Supplied with Enzyme:
10X NEBuffer for Protein Kinases

Reaction Conditions: 1X NEBuffer for Protein Kinases, supplemented with 200 μ M ATP and gamma-labeled ATP to a final specific activity of 100–500 μ Ci/ μ mol. **Incubate at 30°C.**

1X NEBuffer for Protein Kinases:

50 mM Tris-HCl
10 mM MgCl₂
0.1 mM EDTA
2 mM DTT
0.01% Brij 35
pH 7.5 @ 25°C

Note that optimal incubation times and enzyme concentrations must be determined empirically for each particular substrate.

Unit Definition: One unit is defined as the amount of GSK-3 required to catalyze the transfer of 1 pmol of phosphate to CREB Phosphopeptide, KRREILSRPPpSYR (400 μ M, NEB #P6041), in 1 minute at 30°C in a total reaction volume of 25 μ l.

Specific Activity: ~ 5,000,000 units/mg.

Molecular Weight: 47 kDa.

Quality Assurance: GSK-3 contains no detectable protease or phosphatase activities.

Quality Control Assays

Protease Activity: After incubation of 1,000 units of Glycogen Synthase Kinase 3 (GSK-3) with 0.2 nmol of a standard mixture of proteins for 2 hours at 30°C, no proteolytic activity could be detected by SDS-PAGE analysis.

Phosphatase Activity: After incubation of 1,000 units of Glycogen Synthase Kinase 3 (GSK-3) with 50 mM *p*-nitrophenyl phosphate for 2 hours at 30°C, no phosphatase activity could be detected by spectrophotometric analysis.

References:

1. Embi, N., Rylatt, D.B. and Cohen, P. (1980) *Eur. J. Biochem.* 107, 519–527.
2. Hemmings, B.A. et al. (1982) *Eur. J. Biochem.* 119, 443–451.
3. Vandenhede, J.R. et al. (1980) *J. Biol. Chem.* 255, 11768–11774.

(See other side)

CERTIFICATE OF ANALYSIS

Glycogen Synthase Kinase 3 (GSK-3)



1-800-632-7799
info@neb.com
www.neb.com



P6040S 029160217021

P6040S



10,000 units 500,000 U/ml Lot: 0291602
RECOMBINANT Store at -20°C Exp: 2/17

Description: Glycogen Synthase Kinase 3 (GSK-3) is a serine/threonine protein kinase and one of several protein kinases, which phosphorylate glycogen synthase. It is also called Factor A (F_a) for its ability to activate the MgATP-dependent form of the protein phosphatase PP1 called F_c (1–4). Recent studies demonstrate that GSK-3 can autophosphorylate Ser, Thr and Tyr. Ser/Thr phosphorylation causes inactivation, and

New Reaction Buffer

Molecular Weight: 47 kDa.

Quality Assurance: GSK-3 contains no detectable protease or phosphatase activities.

Quality Control Assays

Protease Activity: After incubation of 1,000 units of Glycogen Synthase Kinase 3 (GSK-3) with 0.2 nmol of a standard mixture of proteins for 2 hours at 30°C, no proteolytic activity could be detected by SDS-PAGE analysis.

Phosphatase Activity: After incubation of 1,000 units of Glycogen Synthase Kinase 3 (GSK-3) with 50 mM *p*-nitrophenyl phosphate for 2 hours at 30°C, no phosphatase activity could be detected by spectrophotometric analysis.

References:

1. Embi, N., Rylatt, D.B. and Cohen, P. (1980) *Eur. J. Biochem.* 107, 519–527.
2. Hemmings, B.A. et al. (1982) *Eur. J. Biochem.* 119, 443–451.
3. Vandenhede, J.R. et al. (1980) *J. Biol. Chem.* 255, 11768–11774.

(See other side)

CERTIFICATE OF ANALYSIS

4. Woodgett, J.R. (1990) *EMBO J.* 9, 2431–2438.
5. Wang, Q.M. et al. (1994) *J. Biol. Chem.* 269, 14566–14574.
6. Cole, A. et al. (2004) *Biochem. J.* 377, 249–255.
7. Frame, S. and Cohen P. (2001) *Biochem. J.* 359, 1–16.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Page 2 (P6040)

4. Woodgett, J.R. (1990) *EMBO J.* 9, 2431–2438.
5. Wang, Q.M. et al. (1994) *J. Biol. Chem.* 269, 14566–14574.
6. Cole, A. et al. (2004) *Biochem. J.* 377, 249–255.
7. Frame, S. and Cohen P. (2001) *Biochem. J.* 359, 1–16.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Page 2 (P6040)