

Enterokinase, light chain



1-800-632-7799
info@neb.com
www.neb.com



P8070S 041150716071

P8070S



0.063 µg **2.0 µg/ml** **Lot: 0411507**
RECOMBINANT **Store at -20°C** **Exp: 7/16**

Description: Enterokinase is a specific protease that cleaves after lysine at its cleavage site, Asp-Asp-Asp-Asp-Lys. It will sometimes cleave at other basic residues, depending on the conformation of the protein substrate. Enterokinase will not cleave a site followed by proline.

Note: Now sold by weight

Enterokinase, light chain



1-800-632-7799
info@neb.com
www.neb.com



P8070S 041150716071

P8070S



0.063 µg **2.0 µg/ml** **Lot: 0411507**
RECOMBINANT **Store at -20°C** **Exp: 7/16**

Description: Enterokinase is a specific protease that cleaves after lysine at its cleavage site, Asp-Asp-Asp-Asp-Lys. It will sometimes cleave at other basic residues, depending on the conformation of the protein substrate. Enterokinase will not cleave a site followed by proline.

Note: Now sold by weight

Source: This preparation is purified from *K. lactis* containing a clone of the light chain of the bovine enterokinase gene (1,2).

Supplied in: 20 mM Tris-HCl, 200 mM NaCl, 2 mM CaCl₂, 50% glycerol (pH 7.2 @ 4°C).

Molecular Weight: 26.3 kDa. Its apparent molecular weight on SDS-PAGE is 31 kDa.

Suggested Reaction Conditions: The amount of enzyme required to cleave a fusion protein in a 16 hour reaction at room temperature ranges from 0.0001% to 0.5% (w/w). Cleavage of an MPB-paramyosin-ΔSal fusion protein with an enterokinase site requires 0.0006%.

Unit Definition: 0.00016 µg of Enterokinase will cleave 25 µg of test substrate to 95% completion in 16 hours or less at 25°C.

Unit Assay Conditions: 20 mM Tris-HCl (pH 8.0 @ 25°C), 50 mM NaCl, 2 mM CaCl₂, 25 µg of an MBP fusion protein test substrate and enzyme.
Incubate at 23°C

Source: This preparation is purified from *K. lactis* containing a clone of the light chain of the bovine enterokinase gene (1,2).

Supplied in: 20 mM Tris-HCl, 200 mM NaCl, 2 mM CaCl₂, 50% glycerol (pH 7.2 @ 4°C).

Molecular Weight: 26.3 kDa. Its apparent molecular weight on SDS-PAGE is 31 kDa.

Suggested Reaction Conditions: The amount of enzyme required to cleave a fusion protein in a 16 hour reaction at room temperature ranges from 0.0001% to 0.5% (w/w). Cleavage of an MPB-paramyosin-ΔSal fusion protein with an enterokinase site requires 0.0006%.

Unit Definition: 0.00016 µg of Enterokinase will cleave 25 µg of test substrate to 95% completion in 16 hours or less at 25°C.

Unit Assay Conditions: 20 mM Tris-HCl (pH 8.0 @ 25°C), 50 mM NaCl, 2 mM CaCl₂, 25 µg of an MBP fusion protein test substrate and enzyme.
Incubate at 23°C

Removal: Enterokinase will bind specifically to trypsin inhibitor agarose (e.g., Sigma T-0637).

References:

- Collins-Racie, L.A. et al. (1995) *Biotechnology* 13, 982-987.
- Taron, C. and Colussi, P., unpublished observations.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

Removal: Enterokinase will bind specifically to trypsin inhibitor agarose (e.g., Sigma T-0637).

References:

- Collins-Racie, L.A. et al. (1995) *Biotechnology* 13, 982-987.
- Taron, C. and Colussi, P., unpublished observations.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS