

m⁷G(5')ppp(5')A RNA Cap Structure Analog



1-800-632-7799
info@neb.com
www.neb.com



S1405S 001130516051

S1405S

1 μ mol Lot: 0011305

Store at -20°C Exp: 5/16

m⁷G(5')ppp(5')A Sodium Salt

Description: The 5' terminal m⁷G cap present on most eukaryotic mRNAs promotes translation *in vitro* at the initiation level (1,2,3). For most RNAs, elimination of the cap structure causes a loss of stability, especially against exonuclease degradation (4), and a decrease in the formation of the initiation complex of mRNAs for protein synthesis (4,5). Certain prokaryotic mRNAs containing a 5'

terminal cap structure are translated as efficiently as or more efficiently than eukaryotic mRNAs in a eukaryotic cell-free protein synthesizing system (5). Also a cap requirement has been observed for splicing eukaryotic substrate RNAs (6).

A method using *E. coli* RNA polymerase primed with m⁷G(5')ppp(5')G or m⁷G(5')ppp(5')A for an efficient *in vitro* synthesis of capped RNAs has been developed by Contreas (7). Larger amounts of capped RNAs are produced by transcription systems using SP6 RNA polymerase primed with m⁷G(5')ppp(5')G (6).

Quality Controls

The purity and identity of m⁷G(5')ppp(5')A (Cap Analog) is \geq 95% as determined by HPLC analysis and mass spec respectively.

The RNA Cap Structure Analog is functionally tested for recognition by an RNA Polymerase and its incorporation into a run-off transcript.

Molecular Formula: C₂₁H₃₀N₁₀O₁₇P₃ (Free Acid)

Molecular Weight: 787.44 g/mol (Free acid)

Extinction Coefficient: $\lambda_{260} = \sim 22,700$ Lmol⁻¹ cm⁻¹

Note: Addition of 100 μ l water gives approximately a 10 mM solution.

References:

1. Shatkin, A.J. (1978) *Cell* 9, 645–653.
2. Fillipowicz, W. (1978) *FEBS Lett.* 96, 1–11.
3. Banerjee, A.K. (1980) *Microbiol. Rev.* 44, 175–205.
4. Miura, K. (1981) *Adv. Biophys.* 14, 205–238.
5. Shatkin, A.J. et al. (1977) *Nucleic Acids. Res.* 4, 3065–3081.
6. Konarska, M.M. et al. (1984) *Cell* 38, 731–736.
7. Contreas, R. et al. (1982) *Nucleic Acids. Res.* 10, 6353–6363.
8. Paterson, B.M. and Rosenberg, M. (1979) *Nature* 279, 696–701.

CERTIFICATE OF ANALYSIS

m⁷G(5')ppp(5')A RNA Cap Structure Analog



1-800-632-7799
info@neb.com
www.neb.com



S1405S 001130516051

S1405S

1 μ mol Lot: 0011305

Store at -20°C Exp: 5/16

m⁷G(5')ppp(5')A Sodium Salt

Description: The 5' terminal m⁷G cap present on most eukaryotic mRNAs promotes translation *in vitro* at the initiation level (1,2,3). For most RNAs, elimination of the cap structure causes a loss of stability, especially against exonuclease degradation (4), and a decrease in the formation of the initiation complex of mRNAs for protein synthesis (4,5). Certain prokaryotic mRNAs containing a 5'

terminal cap structure are translated as efficiently as or more efficiently than eukaryotic mRNAs in a eukaryotic cell-free protein synthesizing system (5). Also a cap requirement has been observed for splicing eukaryotic substrate RNAs (6).

A method using *E. coli* RNA polymerase primed with m⁷G(5')ppp(5')G or m⁷G(5')ppp(5')A for an efficient *in vitro* synthesis of capped RNAs has been developed by Contreas (7). Larger amounts of capped RNAs are produced by transcription systems using SP6 RNA polymerase primed with m⁷G(5')ppp(5')G (6).

Quality Controls

The purity and identity of m⁷G(5')ppp(5')A (Cap Analog) is \geq 95% as determined by HPLC analysis and mass spec respectively.

The RNA Cap Structure Analog is functionally tested for recognition by an RNA Polymerase and its incorporation into a run-off transcript.

Molecular Formula: C₂₁H₃₀N₁₀O₁₇P₃ (Free Acid)

Molecular Weight: 787.44 g/mol (Free acid)

Extinction Coefficient: $\lambda_{260} = \sim 22,700$ Lmol⁻¹ cm⁻¹

Note: Addition of 100 μ l water gives approximately a 10 mM solution.

References:

1. Shatkin, A.J. (1978) *Cell* 9, 645–653.
2. Fillipowicz, W. (1978) *FEBS Lett.* 96, 1–11.
3. Banerjee, A.K. (1980) *Microbiol. Rev.* 44, 175–205.
4. Miura, K. (1981) *Adv. Biophys.* 14, 205–238.
5. Shatkin, A.J. et al. (1977) *Nucleic Acids. Res.* 4, 3065–3081.
6. Konarska, M.M. et al. (1984) *Cell* 38, 731–736.
7. Contreas, R. et al. (1982) *Nucleic Acids. Res.* 10, 6353–6363.
8. Paterson, B.M. and Rosenberg, M. (1979) *Nature* 279, 696–701.

CERTIFICATE OF ANALYSIS