

## New England Biolabs Certificate of Analysis


Product Name: Magnesium Sulfate (MgSO<sub>4</sub>) Solution  
 Catalog Number: B1003S  
 Concentration: 100 mM  
 Lot Number: 10055849  
 Expiration Date: 02/2024  
 Storage Temperature: -20°C  
 Specification Version: PS-B1003S v1.0  
 Composition (1X): 100 mM MgSO<sub>4</sub>

Magnesium Sulfate (MgSO <sub>4</sub> ) Solution Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B1003SVIAL	Magnesium Sulfate (MgSO <sub>4</sub> ) Solution	10042724	Pass

Assay Name/Specification	Lot # 10055849
<p><b>RNase Activity (Extended Digestion)</b>            A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Magnesium Sulfate (MgSO<sub>4</sub>) Solution is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p><b>qPCR DNA Contamination (E. coli Genomic)</b>            A minimum of 1 µl of Magnesium Sulfate (MgSO<sub>4</sub>) Solution is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p><b>Phosphatase Activity (pNPP, Buffer)</b>            A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl<sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 20 µl Magnesium Sulfate (MgSO<sub>4</sub>) Solution incubated for 4 hours at 37°C yields &lt;0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p><b>pH (buffers/solutions)</b>            The pH of 100 mM Magnesium Sulfate (MgSO<sub>4</sub>) Solution is between pH 5.3 and 5.7 at 25°C.</p>	Pass

Assay Name/Specification	Lot # 10055849
<p><b>PCR Amplification (5.0 kb Lambda DNA, Mg2+)</b> A 50 µl reaction in ThermoPol II® (Mg-free) Reaction Buffer containing 2 mM Magnesium Sulfate (MgSO<sub>4</sub>) Solution in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.</p>	<b>Pass</b>
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 µl of Magnesium Sulfate (MgSO<sub>4</sub>) Solution incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 µl of Magnesium Sulfate (MgSO<sub>4</sub>) Solution incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Conductivity (buffers/solutions)</b> The conductivity of 100 mM Magnesium Sulfate (MgSO<sub>4</sub>) Solution is between 8.5 and 10.5 mS/cm at 25°C.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.



Doreen Duquette  
Production Scientist  
23 Apr 2019



Michael Tonello  
Packaging Quality Control Inspector  
25 Sep 2019