

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	NEBufferTM 2
Catalog #:	<i>B7002S</i>
Concentration:	10X Concentrate
Lot #:	0371803
Assay Date:	03/2018
Expiration Date:	03/2021
Storage Temp:	-20°C
Composition (1X):	50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, (pH 7.9 @ 25°C)
Specification Version:	PS-B7002S v1.0
Effective Date:	30 Jan 2018

Assay Name/Specification (minimum release criteria)	Lot #0371803
Conductivity (buffers/solutions) - The conductivity of 10X NEBuffer 2 is between 52 and 78 mS/cm at 25°C.	Pass
Endonuclease Activity (Nicking, Buffer) - A 50 μ l reaction in 1X NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, BSA, Buffer) - A 50 μ l reaction in 1X NEBuffer 2 plus 100 μ g/ml Bovine Serum Albumin containing 1 μ g of Lambda DNA and 1 unit of Sphl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, BSA, Buffer) - A 50 μ l reaction in 1X NEBuffer 2 plus 100 μ g/ml Bovine Serum Albumin containing 1 μ g of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) - A 50 μ l reaction in 1X NEBuffer 2 containing 1 μ g of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
pH (buffers/solutions) - The pH of 10X NEBuffer 2 is between pH 7.8 and 8.0 at 25°C.	Pass
RNase Activity (Buffer) - A 10 µl reaction in 1X NEBuffer 2 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass

otunen

Authorized by Derek Robinson 30 Jan 2018



(Uh Int

Inspected by Tony Spear-Alfonso 23 Mar 2018