

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	NEBuffer™ 2
Catalog Number:	B7002S
Concentration:	10 X Concentrate
Lot Number:	10018177
Expiration Date:	06/2021
Storage Temperature:	-20°C
Specification Version:	PS-B7002S v1.0
Composition (1X):	50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, (pH 7.9 @ 25°C)

NEBuffer™ 2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B7002SVIAL	NEBuffer™ 2	10013286	Pass	

Assay Name/Specification	Lot # 10018177
RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer 2 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass
pH (buffers/solutions) The pH of 10X NEBuffer 2 is between pH 7.8 and 8.0 at 25°C.	Pass
Conductivity (buffers/solutions) The conductivity of 10X NEBuffer 2 is between 52 and 78 mS/cm at 25°C.	Pass
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, BSA, Buffer) A 50 μl reaction in 1X NEBuffer 2 plus 100 μg/ml Bovine Serum Albumin containing 1 μg of Lambda DNA and 1 unit of Sphl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, BSA, Buffer) A 50 μl reaction in 1X NEBuffer 2 plus 100 μg/ml Bovine Serum Albumin containing 1	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10018177
µg of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Go

Tony Spear-Alfonso Production Scientist 28 Jun 2018

on. W. Michae

Michael Tonello Packaging Quality Control Inspector 03 Aug 2018

