

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: NEBuffer™ 2

Catalog Number: B7002S

Concentration: 10 X Concentrate

Lot Number: 10034583
Expiration Date: 01/2022
Storage Temperature: -20°C

Specification Version: PS-B7002S v1.0

Composition (1X): 50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, (pH 7.9 @ 25°C)

NEBuffer™ 2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B7002SVIAL	NEBuffer™ 2	10033147	Pass	

Assay Name/Specification	Lot # 10034583
Conductivity (buffers/solutions)	Pass
The conductivity of 10X NEBuffer 2 is between 52 and 78 mS/cm at 25°C.	
Endonuclease Activity (Nicking, Buffer)	Pass
A 50 μl reaction in 1X NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as	
determined by agarose gel electrophoresis.	
	_
Functional Testing (Restriction Digest, BSA, Buffer) A 50 µl reaction in 1X NEBuffer 2 plus 100 µg/ml Bovine Serum Albumin containing 1	Pass
μg of Lambda DNA and 1 unit of Sphl incubated for 1 hour at 37°C results in complete	
digestion of the substrate DNA as determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, BSA, Buffer)	Pass
A 50 μl reaction in 1X NEBuffer 2 plus 100 μg/ml Bovine Serum Albumin containing 1	
μg of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel	
electrophoresis.	
Non-Specific DNase Activity (16 hour, Buffer)	Pass
A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of PhiX174-HaeIII DNA incubated	1 433
for 16 hours at 37°C results in a DNA pattern free of detectable nuclease	
degradation as determined by agarose gel electrophoresis.	



B7002S / Lot: 10034583

Page 1 of 2

Assay Name/Specification	Lot # 10034583
pH (buffers/solutions) The pH of 10X NEBuffer 2 is between pH 7.8 and 8.0 at 25°C.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer 2 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso Production Scientist

NEW ENGLAND

10 Jan 2019

Michael Tonello

Packaging Quality Control Inspector

14 Jan 2019



B7002S / Lot: 10034583

Page 2 of 2