

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	NEBuffer™ 2
Catalog Number:	B7002S
Concentration:	10 X Concentrate
Lot Number:	10043520
Expiration Date:	01/2022
Storage Temperature:	-20°C
Specification Version:	PS-B7002S v1.0
Composition (1X):	50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, (pH 7.9 @ 25°C)

NEBuffer™ 2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B7002SVIAL	NEBuffer™ 2	10033147	Pass	

Assay Name/Specification	Lot # 10043520
Conductivity (buffers/solutions)	Pass
The conductivity of 10X NEBuffer 2 is between 52 and 78 mS/cm at 25°C.	
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, BSA, Buffer) A 50 μl reaction in 1X NEBuffer 2 plus 100 μg/ml Bovine Serum Albumin containing 1 μg of Lambda DNA and 1 unit of SphI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, BSA, Buffer) A 50 μ l reaction in 1X NEBuffer 2 plus 100 μ g/ml Bovine Serum Albumin containing 1 μ g of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10043520
pH (buffers/solutions) The pH of 10X NEBuffer 2 is between pH 7.8 and 8.0 at 25°C.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer 2 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Mi

Tony Spear-Alfonso Production Scientist 10 Jan 2019

Michae on l

Michael Tonello Packaging Quality Control Inspector 13 May 2019

