

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: NEBuffer™ 2

Catalog Number: B7002S

Concentration: 10 X Concentrate

Lot Number: 10045727
Expiration Date: 04/2022
Storage Temperature: -20°C

Specification Version: PS-B7002S v1.0

Composition (1X): 50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, (pH 7.9 @ 25°C)

NEBuffer™ 2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B7002SVIAL	NEBuffer™ 2	10042968	Pass	

Assay Name/Specification	Lot # 10045727
Conductivity (buffers/solutions)	Pass
The conductivity of 10X NEBuffer 2 is between 52 and 78 mS/cm at 25°C.	
Endonuclease Activity (Nicking, Buffer)	Pass
A 50 μl reaction in 1X NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA	
incubated for 4 hours at 37°C results in <10% conversion to the nicked form as	
determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, BSA, Buffer)	Pass
A 50 μl reaction in 1X NEBuffer 2 plus 100 μg/ml Bovine Serum Albumin containing 1	
μg of Lambda DNA and 1 unit of Sphl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	
a againete ger encourance 2 in rate determined by againete ger encouragnment en	
Functional Testing (Restriction Digest, BSA, Buffer)	Pass
A 50 μl reaction in 1X NEBuffer 2 plus 100 μg/ml Bovine Serum Albumin containing 1 μg of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in	
complete digestion of the substrate DNA as determined by agarose gel	
electrophoresis.	
Non-Specific DNaso Activity (16 hour Buffer)	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of PhiX174-HaeIII DNA incubated	F455
for 16 hours at 37°C results in a DNA pattern free of detectable nuclease	
degradation as determined by agarose gel electrophoresis.	



B7002S / Lot: 10045727

Page 1 of 2

Assay Name/Specification	Lot # 10045727
pH (buffers/solutions) The pH of 10X NEBuffer 2 is between pH 7.8 and 8.0 at 25°C.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer 2 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Production Scientist

24 Apr 2019

Michael Tonello

Packaging Quality Control Inspector

28 Jun 2019



B7002S / Lot: 10045727

Page 2 of 2