

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	NEBuffer™ 2
Catalog Number:	B7002S
Concentration:	10 X Concentrate
Packaging Lot Number:	10069218
Expiration Date:	04/2023
Storage Temperature:	-20°C
Specification Version:	PS-B7002S v1.0
Composition (1X):	50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, (pH 7.9 @ 25°C)

NEBuffer™ 2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B7002SVIAL	NEBuffer™ 2	10074105	Pass	

Assay Name/Specification	Lot # 10069218
Conductivity (buffers/solutions)	Pass
The conductivity of 10X NEBuffer 2 is between 52 and 78 mS/cm at 25°C.	
Endenuelesse Astivity (Nicking Buffer)	Pass
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA	Fass
incubated for 4 hours at 37°C results in <10% conversion to the nicked form as	
determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, BSA, Buffer)	Pass
A 50 µl reaction in 1X NEBuffer 2 plus 100 µg/ml Bovine Serum Albumin containing 1	
µg of Lambda DNA and 1 unit of Sphl incubated for 1 hour at 37°C results in complete	
digestion of the substrate DNA as determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, BSA, Buffer)	Pass
A 50 µl reaction in 1X NEBuffer 2 plus 100 µg/ml Bovine Serum Albumin containing 1	
$\mu$ g of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in	
complete digestion of the substrate DNA as determined by agarose gel	
electrophoresis.	
Non-Specific DNase Activity (16 hour, Buffer)	Pass
A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of PhiX174-HaeIII DNA incubated	
for 16 hours at 37°C results in a DNA pattern free of detectable nuclease	
degradation as determined by agarose gel electrophoresis.	





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10069218
<b>pH (buffers/solutions)</b> The pH of 10X NEBuffer 2 is between pH 7.8 and 8.0 at 25°C.	Pass
<b>RNase Activity (Buffer)</b> A 10 μl reaction in 1X NEBuffer 2 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Michael Dalton Production Scientist 29 Apr 2020

val-

Av Minichiello Packaging Quality Control Inspector 29 Apr 2020

