

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	NEBuffer™ 4
Catalog Number:	B7004S
Concentration:	10 X Concentrate
Lot Number:	10009147
Expiration Date:	10/2020
Storage Temperature:	-20°C
Specification Version:	PS-B7004S v1.0
Composition (1X):	50 mM Potassium Acetate , 20 mM Tris-acetate, 10 mM Magnesium Acetate, 1 mM DTT, (pH 7.9 @ 25°C)

NEBuffer™ 4 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B7004SVIAL	NEBuffer™ 4	0181710	Pass	

Assay Name/Specification	Lot # 10009147
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBuffer 4 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
pH (buffers/solutions) The pH of 10X NEBuffer 4 is between pH 7.8 and 8.0 at 25°C.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass
Endonuclease Activity (Nicking, Buffer) A 50 μ I reaction in 1X NEBuffer 4 containing 1 μ g of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Conductivity (buffers/solutions) The conductivity of 10X NEBuffer 4 is between 36 and 54 mS at 25°C.	Pass
Functional Testing (Restriction Digest, BSA, Buffer)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10009147
A 50 μ I reaction in 1X NEBuffer 4 plus 100 μ g/ml Bovine Serum Albumin containing 1 μ g of Lambda dam- DNA and 1 unit of ClaI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, BSA, Buffer) A 50 µl reaction in 1X NEBuffer 4 plus 100 µg/ml Bovine Serum Albumin containing 1 µg of Lambda DNA and 1 unit of Mscl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Michael Dalton Production Scientist 24 May 2018

ton N. Michae

Michael Tonello Packaging Quality Control Inspector 24 May 2018

