

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	CutSmart® Buffer
Catalog Number:	B7204S
Concentration:	10 X Concentrate
Lot Number:	10050604
Expiration Date:	07/2022
Storage Temperature:	-20°C
Specification Version:	PS-B7204S v1.0
Composition (1X):	50 mM Potassium Acetate , 20 mM Tris Acetate, 10 mM Magnesium Acetate, 100 μg/ml BSA, (pH 7.9 @ 25°C)

CutSmart® Buffer Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B7204SVIAL	CutSmart® Buffer	10046080	Pass	

Assay Name/Specification	Lot # 10050604
<b>Functional Testing (Restriction Digest, Buffer)</b> A 50 µl reaction in 1X CutSmart® Buffer containing 1 µg of Lambda DNA and 1 unit of MscI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 μl reaction in 1X CutSmart® Buffer containing 1 μg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>pH (buffers/solutions)</b> The pH of 10X CutSmart® Buffer is between pH 7.8 and 8.0 at 25°C.	Pass
<b>RNase Activity (Buffer)</b> A 10 µl reaction in 1X CutSmart® Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass
<b>Conductivity (buffers/solutions)</b> The conductivity of 10X CutSmart® Buffer is between 40 and 46 mS at 25°C.	Pass
Endonuclease Activity (Nicking, Buffer)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10050604
A 50 µl reaction in 1X CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
<b>Functional Testing (Restriction Digest, Buffer)</b> A 50 µl reaction in 1X CutSmart® Buffer containing 1 µg of Lambda dam- DNA and 1 unit of Clal incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Jean Bastalle

Jaan Bastable Production Scientist 03 Jul 2019

Howellow Michae

Michael Tonello Packaging Quality Control Inspector 18 Jul 2019

