

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | CutSmart® Buffer |
|------------------------|---|
| Catalog #: | B7204S |
| Concentration: | 10X Concentrate |
| <i>Lot</i> #: | 3051803 |
| Assay Date: | 03/2018 |
| Expiration Date: | 03/2021 |
| Storage Temp: | -20°C |
| Composition (1X): | 50 mM Potassium Acetate , 20 mM Tris Acetate, 10 mM Magnesium Acetate, 100 μg/ml BSA, (pH 7.9 @ 25°C) |
| Specification Version: | PS-B7204S v1.0 |
| Effective Date: | 31 Jan 2018 |
| | |

| Assay Name/Specification (minimum release criteria) | Lot #3051803 |
|--|--------------|
| Conductivity (buffers/solutions) - The conductivity of 10X CutSmart® Buffer is between 40 and 46 mS at 25°C. | Pass |
| Endonuclease Activity (Nicking, Buffer) - A 50 μ l reaction in 1X CutSmart® Buffer containing 1 μ g of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Functional Testing (Restriction Digest, Buffer) - A 50 µl reaction in 1X CutSmart® Buffer containing 1 µg of Lambda dam- DNA and 1 unit of ClaI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis. | Pass |
| Functional Testing (Restriction Digest, Buffer) - A 50 µl reaction in 1X CutSmart® Buffer containing 1 µg of Lambda DNA and 1 unit of MscI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in 1X CutSmart® Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| pH (buffers/solutions) - The pH of 10X CutSmart® Buffer is between pH 7.8 and 8.0 at 25°C. | Pass |
| RNase Activity (Buffer) - A 10 μ l reaction in 1X CutSmart® Buffer containing 40 ng of a 300 base single- stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection. | Pass |



B7204S Lot: 3051803 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

timen

Authorized by Derek Robinson 31 Jan 2018



[Un

Inspected by Tony Spear-Alfonso 17 Apr 2018