

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Diluent A
Catalog Number:	B8001S
Concentration:	1 X Concentrate
Packaging Lot Number:	10069211
Expiration Date:	04/2022
Storage Temperature:	-20°C
Specification Version:	PS-B8001S v1.0
Composition (1X):	10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 200 μg/ml BSA , 50 % Glycerol, (pH 7.4 @ 25°C)

Diluent A Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B8001SVIAL	Diluent A	10042697	Pass	

Assay Name/Specification	Lot # 10069211
RNase Activity (Extended Digestion) A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Diluent A is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 1 μ I of Diluent A is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is \leq 1 E. coli genome.	Pass
pH (buffers/solutions) The pH of 1X Diluent A is between pH 7.3 and 7.5 at 25°C.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μ l reaction in CutSmart® Buffer containing 1 μ g of PhiX174-HaeIII DNA and a minimum of 10 μ l of Diluent A incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10069211
A 50 μ I reaction in CutSmart® Buffer containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 10 μ I of Diluent A incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Michael Dalton

Production Scientist 10 Apr 2020

 A_{-}

An Minichiello Packaging Quality Control Inspector 10 Apr 2020

