

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Diluent A   |
|------------------------|---|
| Catalog Number:        | B8001S  |
| Concentration:         | 1 X Concentrate   |
| Packaging Lot Number:  | 10084300  |
| Expiration Date:       | 07/2023   |
| Storage Temperature:   | -20°C   |
| Specification Version: | PS-B8001S v1.0  |
| Composition (1X):      | 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 200 μg/ml BSA<br>, 50 % Glycerol, (pH 7.4 @ 25°C) |

| Diluent A Component List |                       |            |                      |  |
|--------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number          | Component Description | Lot Number | Individual QC Result |  |
| B8001SVIAL               | Diluent A             | 10080020   | Pass                 |  |

| Assay Name/Specification  | Lot # 10084300 |
|---|----------------|
| <b>qPCR DNA Contamination (E. coli Genomic)</b><br>A minimum of 1 $\mu$ I of Diluent A is screened for the presence of E. coli genomic DNA<br>using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results<br>are quantified using a standard curve generated from purified E. coli genomic DNA.<br>The measured level of E. coli genomic DNA contamination is $\leq$ 1 E. coli genome. | Pass           |
| <b>RNase Activity (Extended Digestion)</b><br>A 10 $\mu$ I reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA<br>and a minimum of 1 $\mu$ I of Diluent A is incubated at 37°C. After incubation for 16<br>hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis<br>using fluorescent detection.   | Pass           |
| <b>Endonuclease Activity (Nicking)</b><br>A 50 $\mu$ I reaction in CutSmart® Buffer containing 1 $\mu$ g of supercoiled PhiX174 DNA and<br>a minimum of 10 $\mu$ I of Diluent A incubated for 4 hours at 37°C results in <10%<br>conversion to the nicked form as determined by agarose gel electrophoresis.  | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in CutSmart® Buffer containing 1 µg of PhiX174-HaeIII DNA and a<br>minimum of 10 µl of Diluent A incubated for 16 hours at 37°C results in a DNA<br>pattern free of detectable nuclease degradation as determined by agarose gel<br>electrophoresis.   | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification                                  | Lot # 10084300 |
|---|----------------|
| pH (buffers/solutions)                                    | Pass           |
| The pH of 1X Diluent A is between pH 7.3 and 7.5 at 25°C. |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Michael Dalton

Michael Dalton Production Scientist 30 Oct 2020

Michae m. 11

Michael Tonello Packaging Quality Control Inspector 30 Oct 2020

