

New England Biolabs Certificate of Analysis

Product Name: Standard Taq (Mg-free) Reaction Buffer Pack
Catalog Number: B9015S
Concentration: 10 X Concentrate
Packaging Lot Number: 10240901
Expiration Date: 07/2028
Storage Temperature: -20°C
Specification Version: PS-B9015S v2.0
Composition (1X): 10 mM Tris-HCl, 50 mM KCl, (pH 8.3 @ 25°C)


Standard Taq (Mg-free) Reaction Buffer Pack Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B9021SVIAL	Magnesium Chloride (MgCl ₂) Solution	10221498	Pass
B9015SVIAL	Standard Taq (Mg-free) Reaction Buffer Pack	10198988	Pass

Assay Name/Specification	Lot # 10240901
Endonuclease Activity (Nicking, Mg-Free Buffer) A 50 µl reaction in 2X Standard Taq (Mg-free) Reaction Buffer and 3 mM MgCl ₂ containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Mg-Free Buffer) A 50 µl reaction in 2X Standard Taq (Mg-free) Reaction Buffer and 3 mM MgCl ₂ containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (5 kb Lambda DNA, Mg-Free Buffer) A 50 µl reaction in Standard Taq (Mg-free) Reaction Buffer and 1.5 mM MgCl ₂ in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.	Pass
Phosphatase Activity (pNPP, Buffer) A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl ₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl Standard Taq (Mg-free) Reaction Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass

Assay Name/Specification	Lot # 10240901
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Standard Taq (Mg-free) Reaction Buffer is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>pH (buffers/solutions) The pH of 10X Standard Taq (Mg-free) Reaction Buffer is between pH 8.2 and 8.4 at 25°C.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic, Buffer) A minimum of 1 µl of Standard Taq (Mg-free) Reaction Buffer is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Lea Antonopoulos
Production Scientist
18 Sep 2023



Talia Monkiewicz
Packaging Quality Control Inspector
11 Jul 2024