

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | NEB® 5-alpha Competent E. coli (High Efficiency) |
|------------------------|--|
| Catalog Number:        | C2987R   |
| Packaging Lot Number:  | 10172451   |
| Expiration Date:       | 10/2024  |
| Storage Temperature:   | -80°C  |
| Specification Version: | PS-C2987R v3.0                                   |

| NEB® 5-alpha Competent E. coli (High Efficiency) Component List |  |            |                      |  |
|---|--|------------|----------------------|--|
| NEB Part Number   | Component Description                            | Lot Number | Individual QC Result |  |
| N3041AVIAL  | pUC19 Vector                                     | 10165704   | Pass                 |  |
| C2987RVIAL  | NEB® 5-alpha Competent E. coli (High Efficiency) | 10168770   | Pass                 |  |
| B9040AVIAL  | Outgrowth Medium 1.5                             | 10168310   | Pass                 |  |

| Assay Name/Specification  | Lot # 10172451 |
|---|----------------|
| Antibiotic Sensitivity (Tetracycline)<br>15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked<br>onto a Rich Broth plate containing Tetracycline will not form colonies after<br>incubation for 16 hours at 37°C.     | Pass           |
| Antibiotic Sensitivity (Nitrofurantoin)<br>15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked<br>onto a Rich Broth plate containing Nitrofurantoin will not form colonies after<br>incubation for 16 hours at 37°C. | Pass           |
| Antibiotic Sensitivity (Kanamycin)<br>15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked<br>onto a Rich Broth plate containing Kanamycin will not form colonies after incubation<br>for 16 hours at 37°C.           | Pass           |
| Antibiotic Sensitivity (Streptomycin)<br>15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked<br>onto a Rich Broth plate containing Streptomycin will not form colonies after<br>incubation for 16 hours at 37°C.     | Pass           |
| Antibiotic Sensitivity (Spectinomycin)<br>15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked<br>onto a Rich Broth plate containing Spectinomycin will not form colonies after                                       | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification   | Lot # 10172451 |
|--|----------------|
| incubation for 16 hours at 37°C.   |                |
| Antibiotic Sensitivity (Ampicillin)<br>15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked<br>onto a Rich Broth plate containing Ampicillin will not form colonies after<br>incubation for 16 hours at 37°C.  | Pass           |
| <b>Antibiotic Sensitivity (Chloramphenicol)</b><br>15 μl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked<br>onto a Rich Broth plate containing Chloramphenicol will not form colonies after<br>incubation for 16 hours at 37°C.                                 | Pass           |
| <b>Transformation Efficiency</b><br>1 well of NEB® 5-alpha Competent E. coli (High Efficiency) cells were transformed<br>with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation<br>overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10e8 cfu/µg of DNA. | Pass           |
| Blue-White Screening ( $\alpha$ -complementation, Competent Cells)<br>NEB® 5-alpha Competent E. coli (High Efficiency) were shown to be suitable for<br>blue/white screening by $\alpha$ -complementation of the $\beta$ -galactosidase gene using pUC19.                                  | Pass           |
| <b>Phage Resistance (<math>\phi</math> 80)</b><br>15 µl of untransformed NEB® 5-alpha Competent E. coli (High Efficiency) streaked<br>onto a Rich Broth plate does not support plaque formation by phage $\phi$ 80 after<br>incubation for 16 hours at 37°C.                               | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lixin An Production Scientist 13 Oct 2022

N

Nick Privitera Packaging Quality Control Inspector 18 Nov 2022

