

New England Biolabs Certificate of Analysis

Product Name: SHuffle® T7 Express lysY Competent E. coli

 Catalog #:
 C3030J

 Lot #:
 0071803

 Assay Date:
 03/2018

 Expiration Date:
 03/2019

 Storage Temp:
 -80°C

Specification Version: PS-C3030J v1.0 Effective Date: 17 Apr 2017

Assay Name/Specification (minimum release criteria)	Lot #0071803
Antibiotic Resistance (Chloramphenicol) - 15 μl of untransformed Shuffle® T7 Express lysY Competent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Chloramphenicol will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Resistance (Nitrofurantoin) - 15 µl of untransformed Shuffle® T7 Express lysY Competent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Resistance (Spectinomycin) - 15 μl of untransformed Shuffle® T7 Express lysY Competent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Spectinomycin will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Resistance (Streptomycin) - 15 µl of untransformed Shuffle® T7 Express lysY Competent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Streptomycin will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Ampicillin) - 15 µl of untransformed Shuffle® T7 Express lysY Competent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Kanamycin) - 15 μl of untransformed Shuffle® T7 Express lysY Competent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Tetracycline) - 15 µl of untransformed Shuffle® T7 Express lysY Competent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.	Pass









New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0071803
Functional Testing (Disulfide Bond Formation) - The nuclease NucA requires disulfide bonds for its stability. When expressed at 37°C in <i>E. coli</i> , NucA is toxic to cells only in its oxidized disulfide-bonded state. Transformation of SHuffle® T7 Express lysY Competent <i>E. coli</i> using 100 pg of plasmid that expresses a MBP-NucA fusion results in < 1% of the colonies when compared to a control transformation of its wild type parent strain NEB 10-beta.	Pass
Phage Resistance (Φ 80) - 15 μ l of untransformed Shuffle® T7 Express lysY Competent <i>E. coli</i> streaked onto a Rich Broth plate does not support plaque formation by phage Φ 80 after incubation for 16 hours at 37°C.	Pass
Transformation Efficiency - 50 μ l of SHuffle® T7 Express lysY Competent <i>E. coli</i> cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10e7 cfu/ μ g of DNA.	Pass

Authorized by Derek Robinson 17 Apr 2017







Inspected by Quiting Ren 18 Apr 2018