

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | SHuffle® T7 Express lysY Competent E.coli |
|------------------------|---|
| Catalog Number:        | C3030J                                    |
| Lot Number:            | 10018139                                  |
| Expiration Date:       | 07/2019                                   |
| Storage Temperature:   | -80°C                                     |
| Specification Version: | PS-C3030J v1.0                            |
| opeonioadon version.   |   |

| SHuffle® T7 Express lysY Competent E.coli Component List |   |            |                      |  |
|--|---|------------|----------------------|--|
| NEB Part Number  | Component Description                     | Lot Number | Individual QC Result |  |
| C3030JVIAL   | SHuffle® T7 Express lysY Competent E.coli | 10015241   | Pass                 |  |

| Assay Name/Specification  | Lot # 10018139 |
|---|----------------|
| <b>Antibiotic Resistance (Chloramphenicol)</b><br>15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB<br>or Rich Broth plate containing Chloramphenicol will form colonies after incubation<br>for 16 hours at 37°C. | Pass           |
| Antibiotic Resistance (Nitrofurantoin)<br>15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB<br>or Rich Broth plate containing Nitrofurantoin will form colonies after incubation<br>for 16 hours at 37°C.          | Pass           |
| Antibiotic Resistance (Spectinomycin)<br>15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB<br>or Rich Broth plate containing Spectinomycin will form colonies after incubation for<br>16 hours at 37°C.            | Pass           |
| Antibiotic Resistance (Streptomycin)<br>15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB<br>or Rich Broth plate containing Streptomycin will form colonies after incubation for<br>16 hours at 37°C.              | Pass           |
| Antibiotic Sensitivity (Ampicillin)<br>15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB<br>or Rich Broth plate containing Ampicillin will not form colonies after incubation<br>for 16 hours at 37°C.             | Pass           |
| Antibiotic Sensitivity (Kanamycin)  | Pass           |





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification  | Lot # 10018139 |
|---|----------------|
| 15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB or Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.   |                |
| Antibiotic Sensitivity (Tetracycline)<br>15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB<br>or Rich Broth plate containing Tetracycline will not form colonies after incubation<br>for 16 hours at 37°C.   | Pass           |
| <b>Functional Testing (Disulfide Bond Formation)</b><br>The nuclease NucA requires disulfide bonds for its stability. When expressed at 37°C in E. coli, NucA is toxic to cells only in its oxidized disulfide-bonded state.<br>Transformation of SHuffle® T7 Express lysY Competent E. coli using 100 pg of plasmid that expresses a MBP-NucA fusion results in < 1% of the colonies when compared to a control transformation of its wild type parent strain NEB 10-beta. | Pass           |
| <b>Phage Resistance (<math>\phi</math> 80)</b><br>15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a<br>Rich Broth plate does not support plaque formation by phage $\phi$ 80 after incubation<br>for 16 hours at 37°C.  | Pass           |
| <b>Transformation Efficiency</b><br>50 $\mu$ I of SHuffle® T7 Express lysY Competent E. coli cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10e7 cfu/ $\mu$ g of DNA.  | Pass           |

This product has been tested and shown to be in compliance with all specifications.

mm

Derek Robinson Production Scientist 16 Jul 2018

Corey Rabeau

Packaging Quality Control Inspector 13 Aug 2018

