

## New England Biolabs Certificate of Analysis

**Product Name:** 5' DNA Adenylation Kit  
**Catalog Number:** E2610S  
**Concentration:** 50 µM  
**Packaging Lot Number:** 10293093  
**Expiration Date:** 05/2027  
**Storage Temperature:** -20°C  
**Specification Version:** PS-E2610S/L v1.0

5' DNA Adenylation Kit Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N0757AVIAL	Adenosine 5' Triphosphate	10289514	Pass
M2611AVIAL	Mth RNA Ligase	10289404	Pass
B2610SVIAL	5' DNA Adenylation Reaction Buffer	10289513	Pass

Assay Name/Specification	Lot # 10293093
<b>* Individual Product Component Note</b> Standard Quality Control Tests are performed for each component included in 5' DNA Adenylation Kit and meet the designated specifications.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in 1X 5' DNA Adenylation Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 pmol of Mth RNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in 1X 5' DNA Adenylation Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 pmol of Mth RNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (Adenylation)</b> A 20 µl reaction in 1X 5' DNA Adenylation Reaction Buffer containing 100 pmol of 17 bp long 5'-ssDNA linker (5'-pCTGTAGGCACCATCAAT-NH <sub>2</sub> -3'), 0.1 mM ATP, and 100 pmol of Mth RNA Ligase incubated for 1 hour at 65°C results in ≥95% 5'-ssDNA linker adenylated.	Pass

Assay Name/Specification	Lot # 10293093
<b>Phosphatase Activity (pNPP)</b> A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 50 pmol of Mth RNA Ligase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> Mth RNA Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>RNase Activity Assay (4 Hour Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 50 pmol of Mth RNA Ligase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Bo Wu  
Production Scientist  
20 May 2025



Talia Monkiewicz  
Packaging Quality Control Inspector  
05 Jun 2025