

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Amylose Resin    |
|------------------------|------------------|
| Catalog Number:        | E8021S           |
| Packaging Lot Number:  | 10064287         |
| Expiration Date:       | 09/2022          |
| Storage Temperature:   | 4°C              |
| Specification Version: | PS-E8021S/L v1.0 |

| Amylose Resin Component List |                       |            |                      |  |
|------------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number              | Component Description | Lot Number | Individual QC Result |  |
| E8021SVIAL                   | Amylose Resin         | 10052501   | Pass                 |  |

| Assay Name/Specification  | Lot # 10064287 |
|---|----------------|
| Functional Binding Assay (Resin Binding Capacity)   | Pass           |
| Amylose Resin (1 ml) was packed into a column and equilibrated with column buffer.          |                |
| Crude extract from E. coli containing a plasmid that expresses a                            |                |
| MBP5*-paramyosin $\Delta$ Sal fusion protein ( 8 ml ) was then passed through the column at |                |
| 25°C, then washed with column buffer and the target protein eluted with 4 ml of             |                |
| column buffer containing 10 mM maltose. Binding capacity was determined to be >4 mg         |                |
| MBP5*-paramyosin $\Delta$ Sal /ml of resin based on A280 of the eluate.                     |                |
| Functional Binding Assay (Resin Binding Specificity)  | Pass           |
| Amylose Resin (1 ml) was packed into a column and equilibrated with column buffer.          | 1 400          |
| Crude extract from E. coli containing a plasmid that expresses a                            |                |
| MBP5*-paramyosin∆Sal fusion protein ( 8 ml ) was then passed through the column at          |                |
| 25°C, and then washed with column buffer. The target protein was eluted with 4 ml of        |                |
| column buffer containing 10 mM maltose. SDS-PAGE of the eluate on a 10-20%                  |                |
| Tris-Glycine gel confirms low non-specific binding of extract proteins and high             |                |
| isolation of target.  |                |

This product has been tested and shown to be in compliance with all specifications.





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Brd

Brad Landgraf Production Scientist 24 Sep 2019

Michae The

Michael Tonello Packaging Quality Control Inspector 23 Jan 2020

