

## New England Biolabs Certificate of Analysis

**Product Name:** T4 Polynucleotide Kinase  
**Catalog Number:** M0201L  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One Richardson unit is defined as the amount of enzyme catalyzing the incorporation of 1 nmol of acid insoluble [<sup>33</sup>P] in 30 minutes at 37°C.  
**Packaging Lot Number:** 10126413  
**Expiration Date:** 04/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 μM ATP, 50 % Glycerol, (pH 7.4 @ 25°C)  
**Specification Version:** PS-M0201S/L v1.0


T4 Polynucleotide Kinase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0201LVIAL	T4 Polynucleotide Kinase	10102421	Pass
B0201SVIAL	T4 Polynucleotide Kinase Reaction Buffer	10119056	Pass

Assay Name/Specification	Lot # 10126413
<p><b>Single Stranded DNase Activity (FAM-Labeled Oligo)</b>            A 50 μl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 50 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C yields &lt;5% degradation as determined by capillary electrophoresis.</p>	Pass
<p><b>qPCR DNA Contamination (E. coli Genomic)</b>            A minimum of 10 units of T4 Polynucleotide Kinase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p><b>RNase Activity (Extended Digestion)</b>            A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of T4 Polynucleotide Kinase is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

Assay Name/Specification	Lot # 10126413
<p><b>Protein Purity Assay (SDS-PAGE)</b> T4 Polynucleotide Kinase is <math>\geq 95\%</math> pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Endonuclease Activity (Nicking)</b> A 50 <math>\mu</math>l reaction in T4 Polynucleotide Kinase Reaction Buffer containing 1 <math>\mu</math>g of supercoiled PhiX174 DNA and a minimum of 100 units of T4 Polynucleotide Kinase incubated for 4 hours at 37°C results in <math>&lt;10\%</math> conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Exonuclease Activity (Radioactivity Release)</b> A 50 <math>\mu</math>l reaction in T4 Polynucleotide Kinase Reaction Buffer containing 1 <math>\mu</math>g of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 100 units of T4 Polynucleotide Kinase incubated for 4 hours at 37°C releases <math>&lt;0.1\%</math> of the total radioactivity.</p>	<b>Pass</b>
<p><b>DNase Activity (Labeled Oligo, 3' extension)</b> A 50 <math>\mu</math>l reaction in CutSmart<sup>®</sup> Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 50 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C yields <math>&lt;5\%</math> degradation as determined by capillary electrophoresis.</p>	<b>Pass</b>
<p><b>DNase Activity (Labeled Oligo, 5' extension)</b> A 50 <math>\mu</math>l reaction in CutSmart<sup>®</sup> Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 50 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C yields <math>&lt;5\%</math> degradation as determined by capillary electrophoresis.</p>	<b>Pass</b>
<p><b>Double Stranded DNase Activity (Labeled Oligo)</b> A 50 <math>\mu</math>l reaction in CutSmart<sup>®</sup> Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 50 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C yields <math>&lt;5\%</math> degradation as determined by capillary electrophoresis.</p>	<b>Pass</b>
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 <math>\mu</math>l reaction in T4 Polynucleotide Kinase Reaction Buffer containing 1 <math>\mu</math>g of Lambda DNA and a minimum of 100 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



---

Ana Egana  
Production Scientist  
05 Nov 2021



---

Michael Tonello  
Packaging Quality Control Inspector  
05 Nov 2021