

New England Biolabs Certificate of Analysis

Product Name: *Klenow Fragment (3'-5' exo-)*
Catalog Number: *M0212S*
Concentration: *5,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 37°C.*
Packaging Lot Number: *10246038*
Expiration Date: *06/2026*
Storage Temperature: *-20°C*
Storage Conditions: *25 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)*
Specification Version: *PS-M0212S/L v2.0*

Klenow Fragment (3'-5' exo-) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0212SVIAL	Klenow Fragment (3'-5' exo-)	10246037	Pass
B7002SVIAL	NEBuffer™ 2	10237091	Pass

Assay Name/Specification	Lot # 10246038
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Klenow Fragment (3'-5' exo-) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 2 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of Klenow Fragment (3'-5' exo-) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 50 units of Klenow Fragment (3'-5' exo-) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Phosphatase Activity (pNPP)	Pass

Assay Name/Specification	Lot # 10246038
<p>A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Klenow Fragment (3'–5' exo-) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	
<p>Protein Purity Assay (SDS-PAGE) Klenow Fragment (3'–5' exo-) is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Klenow Fragment (3'–5' exo-) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescence detection.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in NEBuffer 2 containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 50 units of Klenow Fragment (3'–5' exo-) incubated for 30 minutes at 37°C yields <10% degradation as determined by fluorescent detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 50 units of Klenow Fragment (3'–5' exo-) is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Trinh Nguyen
Production Scientist
18 Jul 2024



Michael Tonello
Packaging Quality Control Inspector
18 Jul 2024