240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: HpaII Methyltransferase

Catalog #: M0214S/L
Concentration: 4,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction

volume of 10 µl against cleavage by HpaII restriction endonuclease.

 Lot #:
 0181609

 Assay Date:
 09/2016

 Expiration Date:
 9/2017

 Storage Temp:
 -20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0214S/L v1.0
Effective Date: 01 Aug 2016


Assay Name/Specification (minimum release criteria)	Lot #0181609
Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in NEBuffer 2.1 containing 1 μg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 40 units of HpaII Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Methylase Activity (dam Methylase) - A 20 μl reaction in HpaII Methyltransferase Buffer supplemented with 80 μM S-adenosylmethionine containing 1 μg Lambda DNA and a minimum of 40 units of HpaII Methyltransferase incubated for 4 hours at 37°C did not protect the DNA from digestion by MboI as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 40 units of HpaII Methyltransferase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) - HpaII Methyltransferase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of HpaII Methyltransferase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

^{*} The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.










Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com



Authorized by Derek Robinson 01 Aug 2016





NEW ENGLAND
BioLabs Inc.



Inspected by
Cathy Shea
21 Sep 2016