

www.neb.com info@neb.com



New England Biolabs Certificate of Analysis

Product Name: Hpall Methyltransferase

Catalog Number: M0214S
Concentration: 4,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl

against cleavage by HpaII restriction endonuclease.

Packaging Lot Number: 10226329
Expiration Date: 12/2024
Storage Temperature: -20°C

Storage Conditions: 150 mM NaCl, 50 mM Tris-HCl, 0.1 mM EDTA, 5 mM TCEP-HCl, 50 %

Glycerol, 200 μg/ml BSA, (pH 7.5 @ 25°C)

Specification Version: PS-M0214S/L v2.0

Hpall Methyltransferase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0214SVIAL	Hpall Methyltransferase	10221815	Pass	
B9003SVIAL	S-adenosylmethionine (SAM)	10210241	Pass	
B6004SVIAL	rCutSmart™ Buffer	10219598	Pass	

Assay Name/Specification	Lot # 10226329
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 40 units of Hpall Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Methylase Activity (dam Methylase) A 20 μl reaction in CutSmart® Buffer supplemented with 80 μM S-adenosylmethionine containing 1 μg Lambda DNA and a minimum of 40 units of Hpall Methyltransferase incubated for 4 hours at 37°C did not protect the DNA from digestion by Mbol as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of HaeIII digested PhiX174 RF I DNA and a minimum of 40 units of HpaII Methyltransferase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



M0214S / Lot: 10226329

Page 1 of 2



Assay Name/Specification	Lot # 10226329
Protein Purity Assay (SDS-PAGE) Hpall Methyltransferase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Hpall Methyltransferase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Nicele Castagnozzi Production Scientist 28 Dec 2023 Michael Tonello

Packaging Quality Control Inspector

10 Jan 2024



M0214S / Lot: 10226329