

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Haelll Methyltransferase

Catalog Number: M0224S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 10 μl

against cleavage by HaeIII restriction endonuclease.

Packaging Lot Number: 10150501
Expiration Date: 05/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM Tris-HCl, 50 mM KCl, 10 mM EDTA, 1 mM DTT, 200 µg/ml BSA, 50 %

Glycerol, (pH 7.5 @ 25°C)

Specification Version: PS-M0224S/L v1.0

HaellI Methyltransferase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0224SVIAL	Haelll Methyltransferase	10150500	Pass	
B9003SVIAL	S-adenosylmethionine (SAM)	10148373	Pass	
B0224SVIAL	HaeIII Methyltransferase Reaction Buffer	10093152	Pass	

Assay Name/Specification	Lot # 10150501
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Haelll Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled pBR322 DNA and a minimum of 100 units of Haelll Methyltransferase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda-HindIII DNA and a minimum of 100 units of Haelll Methyltransferase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



M0224S / Lot: 10150501

Page 1 of 2

Assay Name/Specification	Lot # 10150501
Protein Purity Assay (SDS-PAGE) Haelll Methyltransferase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Haelll Methyltransferase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Timothy Meixsell Production Scientist 18 May 2022 Erin Varney

Packaging Quality Control Inspector

18 May 2022



M0224S / Lot: 10150501

Page 2 of 2