

DISCOVERY 240 County Road
SENUINE Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: 9°N™ DNA Ligase

Catalog Number: M0238S
Concentration: 40,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to give 50%

ligation of the 12-base pair cohesive ends of 1 μg of BstEll-digested Lambda DNA in 15 minutes at 45°C.

Lot Number: 10013497
Expiration Date: 04/2020
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 10 mM (NH4)2SO4, 1 mM DTT, 0.1 mM EDTA,

200 μg/ml BSA, 50 % Glycerol, (pH 7.5 @ 25°C)

Specification Version: PS-M0238S/L v1.0

9°N™ DNA Ligase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0238SVIAL	9°N™ DNA Ligase	10011689	Pass	
B0238SVIAL	10X 9°N™ DNA Ligase Buffer	0011704	Pass	

Assay Name/Specification	Lot # 10013497
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a	Pass
minimum of 400 units of 9°N™ DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in 9°N™ DNA Ligase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 400 units of 9°N™ DNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of Lambda-HindIII DNA and a minimum of 80 units of 9°N™ DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	Pass



M0238S / Lot: 10013497

Page 1 of 2

Assay Name/Specification	Lot # 10013497
and a minimum of 1 µl of 9°N™ DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

Ana Egana
Production Scientist

26 Jun 2018

Michael Tonello

Packaging Quality Control Inspector

26 Jun 2018

