

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: 9°N™ DNA Ligase

Catalog Number: M0238S
Concentration: 40,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to give 50%

ligation of the 12-base pair cohesive ends of 1 μg of BstEll-digested Lambda DNA in 15 minutes at 45°C.

Lot Number: 10053727
Expiration Date: 09/2021
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 10 mM (NH4)2SO4, 1 mM DTT, 0.1 mM EDTA,

200 μg/ml BSA, 50 % Glycerol, (pH 7.5 @ 25°C)

Specification Version: PS-M0238S/L v1.0

9°N™ DNA Ligase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0238SVIAL	9°N™ DNA Ligase	10053726	Pass	
B0238SVIAL	10X 9°N™ DNA Ligase Buffer	10039939	Pass	

Assay Name/Specification	Lot # 10053727
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in 9°N™ DNA Ligase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 400 units of 9°N™ DNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of Lambda-HindIII DNA and a minimum of 80 units of 9°N™ DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of 9°N™ DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Endonuclease Activity (Nicking)	Pass



M0238S / Lot: 10053727

Page 1 of 2



Assay Name/Specification	Lot # 10053727
A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 400 units of 9°N™ DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

1 K brenze

Mary Lorenzen Production Scientist 22 Apr 2019 Josh Hersey

Packaging Quality Control Inspector

19 Sep 2019