

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: 9°N™ DNA Ligase

Catalog Number: M0238S
Concentration: 40,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to give 50%

ligation of the 12-base pair cohesive ends of 1 μg of BstEll-digested Lambda DNA in 15 minutes at 45°C.

Packaging Lot Number: 10064393
Expiration Date: 02/2022
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 10 mM (NH4)2SO4, 1 mM DTT, 0.1 mM EDTA,

200 μg/ml BSA, 50 % Glycerol, (pH 7.5 @ 25°C)

Specification Version: PS-M0238S/L v1.0

| 9°N™ DNA Ligase Component List | | | | |
|--------------------------------|----------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| M0238SVIAL | 9°N™ DNA Ligase | 10064392 | Pass | |
| B0238SVIAL | 10X 9°N™ DNA Ligase Buffer | 10065921 | Pass | |

| Assay Name/Specification | Lot # 10064393 |
|---|----------------|
| Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 4 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 400 units of 9°N™ DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in 9°N™ DNA Ligase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 400 units of 9°N™ DNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of Lambda-HindIII DNA and a minimum of 80 units of 9°N™ DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA | Pass |



M0238S / Lot: 10064393

Page 1 of 2



| Assay Name/Specification | Lot # 10064393 |
|--|----------------|
| and a minimum of 1 µl of 9°N™ DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | |

This product has been tested and shown to be in compliance with all specifications.

Ana Egana

Production Scientist 06 Apr 2020

Jay Minichiello

Packaging Quality Control Inspector

06 Apr 2020

