

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | Micrococcal Nuclease |
|------------------------|---|
| Catalog Number: | M0247S |
| Concentration: | 2,000,000 gel U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in 15 minutes at 37°C, to the extent that the accumulation of low molecular DNA fragments is <400 base pairs as determined by agarose gel electrophoresis. |
| Packaging Lot Number: | 10132870 |
| Expiration Date: | 01/2024 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 50 mM NaCl, 10 mM Tris-HCl, 1 mM EDTA, 50 % Glycerol, (pH 7.5 @ 25°C) |
| Specification Version: | PS-M0247S v1.0 |

| Micrococcal Nuclease Component List | | | | |
|-------------------------------------|-----------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| M0247SVIAL | Micrococcal Nuclease | 10132869 | Pass | |
| B9001SVIAL | Purified BSA | 10119610 | Pass | |
| B0247SVIAL | Micrococcal Nuclease Buffer | 10122127 | Pass | |

| Assay Name/Specification | Lot # 10132870 |
|---|----------------|
| qPCR DNA Contamination (E. coli Genomic) A minimum of 2,000 units of Micrococcal Nuclease is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is \leq 1 E. coli genome. | Pass |
| Protein Purity Assay (SDS-PAGE) Micrococcal Nuclease is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |
| Protease Activity (SDS-PAGE) A 20 μl reaction in 1X Micrococcal Nuclease Reaction Buffer containing 24 μg of a standard mixture of proteins and a minimum of 10,000 units of Micrococcal Nuclease incubated for 16 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection. | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

J. Lorei

John Greci Production Scientist 02 Feb 2022

Michae 11.

Michael Tonello Packaging Quality Control Inspector 02 Feb 2022

