

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: T7 RNA Polymerase

Catalog Number: M0251S
Concentration: 50,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1

nmol ATP into acid-insoluble material in a total reaction volume of 50 µl in 1 hour at 37°C in 1X RNA Polymerase Reaction Buffer.

Lot Number: 10043729
Expiration Date: 04/2021
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl , 50 mM Tris-HCl (pH 7.9), 1 mM EDTA , 20 mM BME , 0.1 %

Triton X-100 , 50 % Glycerol

Specification Version: PS-M0251S/L v3.0

T7 RNA Polymerase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0251SVIAL	T7 RNA Polymerase	10040641	Pass	
B9012SVIAL	RNAPol Reaction Buffer	0181804	Pass	

Assay Name/Specification	Lot # 10043729
Endonuclease Activity (Nicking) A 50 μl reaction in RNAPol Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 150 units of T7 RNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 150 units of T7 RNA Polymerase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 250 units of T7 RNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Promoter Specificity	Pass



M0251S / Lot: 10043729

Page 1 of 2

Assay Name/Specification	Lot # 10043729
A 50 $\mu$ l reaction in RNAPol Reaction Buffer in the presence of 2 mM NTPs containing 1 $\mu$ g of Lambda DNA as a template and a minimum of 200 units of T7 RNA Polymerase incubated for 1 hour at 37°C results in <1.5% of the amount of product incorporated as compared to a control reaction using T7 DNA as a template.	
Protein Purity Assay (SDS-PAGE)  T7 RNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in RNAPol Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 50 units of T7 RNA Polymerase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as	Pass

This product has been tested and shown to be in compliance with all specifications.

determined by gel electrophoresis using fluorescent detection.

Dongxian Yue Production Scientist

22 Apr 2019

Michael Tonello

Packaging Quality Control Inspector

22 Apr 2019

