

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: T7 RNA Polymerase

Catalog Number: M0251S
Concentration: 50,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1

nmol ATP into acid-insoluble material in a total reaction volume of 50 µl in 1 hour at 37°C in 1X RNA Polymerase Reaction Buffer.

Packaging Lot Number: 10091328 Expiration Date: 11/2022 Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl , 50 mM Tris-HCl (pH 7.9), 1 mM EDTA , 20 mM BME , 0.1 %

Triton X-100, 50 % Glycerol

Specification Version: PS-M0251S/L v3.0

| T7 RNA Polymerase Component List |                        |            |                      |  |
|----------------------------------|------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>           | Component Description  | Lot Number | Individual QC Result |  |
| M0251SVIAL                       | T7 RNA Polymerase      | 10091329   | Pass                 |  |
| B9012SVIAL                       | RNAPol Reaction Buffer | 10073286   | Pass                 |  |

| Assay Name/Specification   | Lot # 10091328 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 μl reaction in RNAPol Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 150 units of T7 RNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.              | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 150 units of T7 RNA Polymerase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.                | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 250 units of T7 RNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| Promoter Specificity   | Pass           |



M0251S / Lot: 10091328

Page 1 of 2

| Assay Name/Specification  | Lot # 10091328 |
|---|----------------|
| A 50 µl reaction in RNAPol Reaction Buffer in the presence of 2 mM NTPs containing 1 µg of Lambda DNA as a template and a minimum of 200 units of T7 RNA Polymerase incubated for 1 hour at 37°C results in <1.5% of the amount of product incorporated as compared to a control reaction using T7 DNA as a template.                           |                |
| Protein Purity Assay (SDS-PAGE)  T7 RNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.   | Pass           |
| RNase Activity (Extended Digestion) A 10 µl reaction in RNAPol Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 50 units of T7 RNA Polymerase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist

07 Dec 2020

Michael Tonello

Packaging Quality Control Inspector

07 Dec 2020



M0251S / Lot: 10091328

Page 2 of 2