

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Deep Vent® DNA Polymerase
Catalog #:	M0258S/L
Concentration:	2,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.
Lot #:	0181612
Assay Date:	12/2016
Expiration Date:	12/2018
Storage Temp:	-20°C
Storage Conditions:	10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.1 % Triton®X-100 , 50 % Glycerol, (pH 7.4 @ 25° C)
Specification Version:	PS-M0258S/L v1.0
Effective Date:	22 Apr 2016

Assay Name/Specification (minimum release criteria)	Lot #0181612
Endonuclease Activity (Nicking, Polymerase, dNTP) - A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 400 μ M dNTPs containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 20 units of Deep Vent TM DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
PCR Amplification (2.0 kb Lambda DNA) - A 25 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 5 ng Lambda DNA with 0.5 units of Deep Vent TM DNA Polymerase for 30 cycles of PCR amplification results in the expected 2.0 kb product.	Pass
Phosphatase Activity (pNPP) - A 200 μ l reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 nM <i>p</i> -Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Deep Vent TM DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE) - Deep Vent TM DNA Polymerase is \geq 98% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
qPCR DNA Contamination (<i>E. coli</i> Genomic) - A minimum of 2 units of Deep Vent TM DNA Polymerase is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.	Pass



M0258S/L Lot: 0181612 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0181612
RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single- stranded RNA and a minimum of 1 µl of Deep Vent TM DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

Authorized by Melanie Fortier 22 Apr 2016



UM. Int

Inspected by Tony Spear-Alfonso 01 Dec 2016