240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Therminator<sup>TM</sup> DNA Polymerase

Catalog #: M0261S/L
Concentration: 2,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes

at 75°C.

 Lot #:
 0211803

 Assay Date:
 03/2018

 Expiration Date:
 3/2020

 Storage Temp:
 -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0261S/L v1.0
Effective Date: 30 Jun 2016

Assay Name/Specification (minimum release criteria)	Lot #0211803
<b>Endonuclease Activity (Nicking)</b> - A 50 μl reaction in ThermoPol® Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 units of Therminator <sup>TM</sup> DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 μl reaction in ThermoPol® Reaction Buffer containing 1 μg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 20 units of Therminator <sup>TM</sup> DNA Polymerase incubated for 4 hours at either 37°C or 75°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 2 units of Therminator <sup>TM</sup> DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Phosphatase Activity (pNPP)</b> - A 200 μl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM <i>p</i> -Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Therminator <sup>TM</sup> DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> - Therminator <sup>™</sup> DNA Polymerase is ≥ 98% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0211803
RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-	
stranded RNA and a minimum of 1 µl of Therminator <sup>TM</sup> DNA Polymerase is incubated at 37°C. After incubation	Pass
for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent	Pass
detection.	

M.W. Southworth

Authorized by Maurice Southworth 30 Jun 2016







Inspected by
Tony Spear-Alfonso

02 Mar 2018