

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: phi29 DNA Polymerase

Catalog Number: M0269L Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate

0.5 pmol of dNTP into acid insoluble material in 10 minutes at 30°C.

Lot Number: 10034472
Expiration Date: 11/2020
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.5 % Tween®

20 , 0.5 % IGEPAL® CA-630 , 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0269S/L v2.0

phi29 DNA Polymerase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0269LVIAL	phi29 DNA Polymerase	10032152	Pass	
B9000SVIAL	BSA, Molecular Biology Grade	10028759	Pass	
B0269SVIAL	Ф29 DNA Polymerase Reaction Buffer	10029297	Pass	

Assay Name/Specification	Lot # 10034472
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 100 units of phi29 DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 units of phi29 DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units phi29 DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE)	Pass



M0269L / Lot: 10034472

Page 1 of 2

Assay Name/Specification	Lot # 10034472
phi29 DNA Polymerase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of phi29 DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of phi29 DNA Polymerase is incubated at 37°C. After incubation	Pass

This product has been tested and shown to be in compliance with all specifications.

for 16 hours, >90% of the substrate RNA remains intact as determined by gel

electrophoresis using fluorescent detection.

Tony Spear-Alfonso Production Scientist 14 Nov 2018 Michael Tonello

Packaging Quality Control Inspector

20 Mar 2019



M0269L / Lot: 10034472

Page 2 of 2