

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: E.coli Poly (A) Polymerase

Catalog Number: M0276L
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1

nmol of AMP into RNA in a 20 µl volume in 10 minutes at 37°C.

Packaging Lot Number: 10117867
Expiration Date: 09/2022
Storage Temperature: -20°C

Storage Conditions: 20 mM Tris-HCl, 300 mM NaCl, 1 mM EDTA, 1 mM DTT, 0.1 %

Triton®X-100, 50% Glycerol, (pH 7.5 @ 25°C)

Specification Version: PS-M0276S/L v1.0

E.coli Poly (A) Polymerase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0276LVIAL	E.coli Poly (A) Polymerase	10106644	Pass	
B0756AVIAL	Adenosine-5'-Triphosphate (ATP)	10084290	Pass	
B0276SVIAL	Poly(A) Polymerase Reaction Buffer	10114403	Pass	

Assay Name/Specification	Lot # 10117867
Protein Purity Assay (SDS-PAGE)  E. coli Poly(A) Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using	Pass
Coomassie Blue detection.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in Poly(A) Polymerase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 15 units of E. coli	
Poly(A) Polymerase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
	_
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	Pass
and a minimum of 5 units of E. coli Poly(A) Polymerase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by	
gel electrophoresis using fluorescent detection.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in Poly(A) Polymerase Reaction Buffer containing 1 µg of	



M0276L / Lot: 10117867 Page 1 of 2

Assay Name/Specification	Lot # 10117867
supercoiled PhiX174 DNA and a minimum of 15 units of E. coli Poly(A) Polymerase	
incubated for 4 hours at 37°C results in <10% conversion to the nicked form as	
determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist 16 Aug 2021 Michael Tonello Packaging Quality Control Inspector

16 Aug 2021

