

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Uracil-DNA Glycosylase (UDG)

Catalog Number: M0280L
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the

release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [ <sup>3</sup>H]-uracil in a 50 μl reaction containing 0.2 μg DNA (10<sup>4</sup>-10<sup>5</sup>

cpm/µg) in 30 minutes at 37°C.

Packaging Lot Number: 10087636 Expiration Date: 09/2022 Storage Temperature: -20°C

Storage Conditions: 50 mM KCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol , 100 µg/ml BSA

Specification Version: PS-M0280S/L v1.0

Uracil-DNA Glycosylase (UDG) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0280LVIAL	Uracil-DNA Glycosylase (UDG)	10081988	Pass	
B0280SVIAL	UDG Reaction Buffer	10084484	Pass	

Assay Name/Specification	Lot # 10087636
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 1.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda-HindIII DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose	Pass



M0280L / Lot: 10087636 Page 1 of 2

Assay Name/Specification	Lot # 10087636
gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) Uracil-DNA Glycosylase (UDG) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lauren Higgins Production Scientist 23 Oct 2020

Lauren Higgins

Josh Hersey
Packaging Quality Control Inspector

23 Oct 2020

