

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Uracil-DNA Glycosylase (UDG)
Catalog Number:	M0280S
Concentration:	5,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme that catalyzes the release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [³ H]-uracil in a 50 μl reaction containing 0.2 μg DNA (10 ⁴ -10 ⁵ cpm/μg) in 30 minutes at 37°C.
Lot Number:	10012363
Expiration Date:	06/2020
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 100 μg/ml BSA
Specification Version:	PS-M0280S/L v1.0

Uracil-DNA Glycosylase (UDG) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0280SVIAL	Uracil-DNA Glycosylase (UDG)	10012364	Pass	
B0280SVIAL	UDG Reaction Buffer	0011807	Pass	

Assay Name/Specification	Lot # 10012363
Protein Purity Assay (SDS-PAGE) Uracil-DNA Glycosylase (UDG) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda-HindIII DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10012363
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and	
double-stranded [³ H] E. coli DNA and a minimum of 50 units of Uracil-DNA	
Glycosylase (UDG) incubated for 4 hours at 37°C releases <0.1% of the total	
radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

Lauren Higgins

Lauren Sears Higgins Production Scientist 19 Jun 2018

Non mm

Mary Conflon Packaging Quality Control Inspector 11 Jul 2018

