

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Uracil-DNA Glycosylase (UDG)

Catalog Number: M0280S Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the

release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [³H]-uracil in a 50 μl reaction containing 0.2 μg DNA (10⁴-10⁵

cpm/µg) in 30 minutes at 37°C.

Packaging Lot Number: 10061554
Expiration Date: 07/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, 100 µg/ml BSA

Specification Version: PS-M0280S/L v1.0

Uracil-DNA Glycosylase (UDG) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0280SVIAL	Uracil-DNA Glycosylase (UDG)	10049128	Pass	
B0280SVIAL	UDG Reaction Buffer	10049129	Pass	

Assay Name/Specification	Lot # 10061554
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 1.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda-HindIII DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose	Pass



M0280S / Lot: 10061554

Page 1 of 2

Assay Name/Specification	Lot # 10061554
gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) Uracil-DNA Glycosylase (UDG) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Lauren Sears Higgins Production Scientist

11 Jul 2019

Jay Minichiello

Packaging Quality Control Inspector

19 Dec 2019

