

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Uracil-DNA Glycosylase (UDG)

Catalog Number: M0280S Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the

release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [³H]-uracil in a 50 μl reaction containing 0.2 μg DNA (10⁴-10⁵

cpm/µg) in 30 minutes at 37°C.

Packaging Lot Number: 10100971
Expiration Date: 01/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol , 100 μg/ml BSA

Specification Version: PS-M0280S/L v1.0

Uracil-DNA Glycosylase (UDG) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0280SVIAL	Uracil-DNA Glycosylase (UDG)	10097699	Pass	
B0280SVIAL	UDG Reaction Buffer	10098669	Pass	

Assay Name/Specification	Lot # 10100971
Protein Purity Assay (SDS-PAGE) Uracil-DNA Glycosylase (UDG) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda-HindIII DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 1.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 50 units of Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass



M0280S / Lot: 10100971

Page 1 of 2

Assay Name/Specification	Lot # 10100971
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 50 units of Uracil-DNA	
Glycosylase (UDG) incubated for 4 hours at 37°C releases <0.1% of the total	
radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lauren Higgins Production Scientist 09 Mar 2021

Lauren Higgins

Michael Tonello

Packaging Quality Control Inspector

09 Mar 2021

