

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Topoisomerase I (E. coli)  |
|------------------------|--|
| Catalog Number:        | M0301S   |
| Concentration:         | 5,000 U/ml   |
| Unit Definition:       | One unit is defined as the amount of enzyme that catalyzes the relaxation of > 95% of 0.5 $\mu$ g of negatively supercoiled pUC19 RF I DNA in a total reaction volume of 25 $\mu$ l in 15 minutes at 37°C. |
| Packaging Lot Number:  | 10090025   |
| Expiration Date:       | 09/2021  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 10 mM Tris-HCl, 50 mM KCl, 35 mM (NH4)2SO4, 1 mM DTT, 0.1 mM EDTA,<br>50 % Glycerol, (pH 7.5 @ 25℃)  |
| Specification Version: | PS-M0301S/L v1.0   |

| Topoisomerase I (E. coli) Component List |                           |            |                      |  |
|--|---------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>                   | Component Description     | Lot Number | Individual QC Result |  |
| M0301SVIAL                               | Topoisomerase I (E. coli) | 10081885   | Pass                 |  |
| B7204SVIAL                               | CutSmart® Buffer          | 10085424   | Pass                 |  |

| Assay Name/Specification   | Lot # 10090025 |
|--|----------------|
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and   | Pass           |
| double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 units of Topoisomerase I (E.   |                |
| coli) incubated for 4 hours at 37°C releases <0.5% of the total radioactivity.   |                |
| qPCR DNA Contamination (E. coli Genomic)   | Pass           |
| A minimum of 5 units of Topoisomerase I (E. coli) is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is $\leq$ 1 E. |                |
| coli genome.   |                |
| RNase Activity (Extended Digestion)  | Pass           |
| A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA  |                |
| and a minimum of 1 µl of Topoisomerase I (E. coli) is incubated at 37°C. After   |                |
| incubation for 16 hours, >90% of the substrate RNA remains intact as determined by   |                |
| gel electrophoresis using fluorescent detection.   |                |

This product has been tested and shown to be in compliance with all specifications.





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 03 Nov 2020

Ton. Ol Michae

Michael Tonello Packaging Quality Control Inspector 03 Nov 2020

