

New England Biolabs Certificate of Analysis

Product Name: *RNase Inhibitor, Human Placenta*
Catalog Number: *M0307S*
Concentration: *40,000 U/ml*
Unit Definition: *One unit is defined as the amount of RNase Inhibitor, Human Placenta required to inhibit the activity of 5 ng of RNase A by 50%. Activity is measured by the inhibition of hydrolysis of cytidine 2', 3'-cyclic monophosphate by RNase A.*
Packaging Lot Number: *10173154*
Expiration Date: *09/2024*
Storage Temperature: *-20°C*
Storage Conditions: *50 mM KCl, 20 mM HEPES (pH 7.6), 8 mM DTT, 50 % Glycerol*
Specification Version: *PS-M0307S/L v1.0*

RNase Inhibitor, Human Placenta Component List


NEB Part Number	Component Description	Lot Number	Individual QC Result
M0307SVIAL	RNase Inhibitor, Human Placenta	10169025	Pass

Assay Name/Specification	Lot # 10173154
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 4 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of RNase Inhibitor, Human Placenta incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Latent RNase Activity (Extended Digest) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of heat inactivated RNase Inhibitor, Human Placenta is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 40 units of RNase Inhibitor, Human Placenta incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE)</p>	Pass


Assay Name/Specification	Lot # 10173154
<p>RNase Inhibitor, Human Placenta is $\geq 95\%$ pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p> <p>RNase Activity (Extended Digestion) A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of RNase Inhibitor, Human Placenta is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Dongxian Yue
Production Scientist
21 Nov 2022



Michael Tonello
Packaging Quality Control Inspector
23 Nov 2022