

www.neb.com info@neb.com



## New England Biolabs Certificate of Analysis

Product Name: Human Alkyladenine Glycosylase (hAAG)

Catalog Number: M0313S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to create an AP

site from 1 pmol of a 34-mer oligonucleotide duplex containing a single deoxyinosine site in a total reaction volume of 10  $\mu$ l in 1

hour at 37°C.

Packaging Lot Number: 10279065
Expiration Date: 12/2026
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.5 % Tween®

20 , 0.5 % IGEPAL® CA-630 , 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0313S/L v1.0

| Human Alkyladenine Glycosylase (hAAG) Component List |                                       |            |                      |  |
|--|---------------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>                               | Component Description                 | Lot Number | Individual QC Result |  |
| M0313SVIAL   | Human Alkyladenine Glycosylase (hAAG) | 10260731   | Pass                 |  |
| B9004SVIAL   | ThermoPol® Reaction Buffer Pack       | 10268442   | Pass                 |  |

| Assay Name/Specification  | Lot # 10279065 |
|---|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of hAAG incubated for 4 hours at 37°C results n <10% conversion to the nicked form as determined by agarose gel electrophoresis.                       | Pass           |
| Exonuclease Activity (Radioactivity Release) a 50 µl reaction in NEBuffer 1 containing 1 µg of a mixture of single and louble-stranded [ ³H] E. coli DNA and a minimum of 50 units of hAAG incubated for 4 lours at 37°C releases <0.1% of the total radioactivity.   | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 units of hAAG incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| Protein Purity Assay (SDS-PAGE)   | Pass           |



M0313S / Lot: 10279065

Page 1 of 2



| Assay Name/Specification   | Lot # 10279065 |
|--|----------------|
| hAAG is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue |                |
| detection.   |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Jamie Souza

Production Scientist

19 Dec 2024

Michael Tonello

Packaging Quality Control Inspector

19 Mar 2025



M0313S / Lot: 10279065

Page 2 of 2