

New England Biolabs Certificate of Analysis

Product Name: *RNase Inhibitor, Murine*
Catalog Number: *M0314L*
Concentration: *40,000 U/ml*
Unit Definition: *One unit is defined as the amount of Murine RNase Inhibitor required to inhibit the activity of 5ng of RNase A by 50%. Activity is measured by the inhibition of hydrolysis of cytidine 2', 3'-cyclic monophosphate by RNase A.*
Packaging Lot Number: *10111205*
Expiration Date: *06/2023*
Storage Temperature: *-20°C*
Storage Conditions: *50 mM KCl, 20 mM HEPES (pH 7.6), 8 mM DTT, 50 % Glycerol*
Specification Version: *PS-M0314S/L v1.0*


RNase Inhibitor, Murine Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0314LVIAL	RNase Inhibitor, Murine	10109058	Pass

Assay Name/Specification	Lot # 10111205
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of RNase Inhibitor, Murine is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Protein Purity Assay (SDS-PAGE) RNase Inhibitor, Murine is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Latent RNase Activity (Extended Digest) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of heat inactivated RNase Inhibitor, Murine is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 4 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of RNase Inhibitor,	Pass

Assay Name/Specification	Lot # 10111205
<p>Murine incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p> <p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 40 units of RNase Inhibitor, Murine incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Bhairavi Jani
Production Scientist
23 Jun 2021



Josh Hersey
Packaging Quality Control Inspector
23 Jun 2021